


# Asymptotics with $\Lambda > 0$



Béatrice Bonga, Penn State

*Work in collaboration with A. Asthekar and A. Kesavan  
[PRD 2015 + PRL 2016]*

# Gravitational waves with $\Lambda = 0$



# Gravitational waves with $\Lambda > 0$

---

## Open issues

- ▶ No Bondi news
- ▶ No ‘peeling’ theorem
- ▶ No positivity of energy
- ▶ No positive and negative frequency decomposition needed for Hilbert spaces in quantum theory

# Gravitational waves with $\Lambda > 0$

---

## Open issues

- ▶ No Bondi news
- ▶ No ‘peeling’ theorem
- ▶ No positivity of energy
- ▶ No positive and negative frequency decomposition needed for Hilbert spaces in quantum theory
  
- ▶ ... and no quadrupole formula!

# Outline

---

1. Steps calculation in Minkowski space-time
2. Steps calculation in de Sitter space-time
3. Result

# Linearization

---

One-parameter family of metrics

$$g_{\mu\nu}(\epsilon) = g_{\mu\nu}(0) + \epsilon h_{\mu\nu} + \dots$$

- ▶ For Minkowski:  $g_{\mu\nu}(0) = \eta_{\mu\nu}$
- ▶ For de Sitter:  $g_{\mu\nu}(0) = a^2(\eta) \mathring{g}_{\mu\nu}$  with  $a(\eta) = -\frac{1}{\sqrt{\Lambda/3\eta}}$

# Main ingredients

---

3 main ingredients

1. gauge fixed gravitational wave  $\bar{h}_{\mu\nu}$

$$\square \bar{h}_{\mu\nu} = 16\pi G T_{\mu\nu} \implies \bar{h}_{ij} = \frac{4G}{r} \int d^3x' T_{ij}(t-r, x')$$

2. energy & power

$$H_t = \frac{1}{32\pi G} \int d^3x \left( \dot{h}_{ij} \dot{h}^{ij} - h_{ij} \ddot{h}^{ij} \right)$$

$$P_t := \mathcal{L}_t H_t = \frac{\partial}{\partial t} H_t$$

3. quadrupole moment & conservation of stress-energy tensor

$$Q_{ij} := \int d^3x \rho x_i x_j$$

$$\int d^3x T_{ij} = \frac{1}{2} \int d^3x \tilde{T}_{00} x_i x_j = \frac{1}{2} \ddot{Q}_{ij}$$

# Main ingredients

---

3 main ingredients

1. gauge fixed gravitational wave  $\bar{h}_{\mu\nu}$

$$\square \bar{h}_{\mu\nu} = 16\pi G T_{\mu\nu} \implies \bar{h}_{ij} = \frac{4G}{r} \int d^3x' T_{ij}(t-r, x')$$

2. energy & power

$$H_t = \frac{1}{32\pi G} \int d^3x \left( \dot{h}_{ij} \dot{h}^{ij} - h_{ij} \ddot{h}^{ij} \right)$$

$$P_t := \mathcal{L}_t H_t = \frac{\partial}{\partial t} H_t$$

3. quadrupole moment & conservation of stress-energy tensor

$$Q_{ij} := \int d^3x \rho x_i x_j$$

$$\int d^3x T_{ij} = \frac{1}{2} \int d^3x \tilde{T}_{00} x_i x_j = \frac{1}{2} \ddot{Q}_{ij}$$

# Main ingredients

---

3 main ingredients

1. gauge fixed gravitational wave  $\bar{h}_{\mu\nu}$

$$\square \bar{h}_{\mu\nu} = 16\pi G T_{\mu\nu} \implies \bar{h}_{ij} = \frac{4G}{r} \int d^3x' T_{ij}(t-r, x')$$

2. energy & power

$$H_t = \frac{1}{32\pi G} \int d^3x \left( \dot{h}_{ij} \dot{h}^{ij} - h_{ij} \ddot{h}^{ij} \right)$$

$$P_t := \mathcal{L}_t H_t = \frac{\partial}{\partial t} H_t$$

3. quadrupole moment & conservation of stress-energy tensor

$$Q_{ij} := \int d^3x \rho x_i x_j$$

$$\int d^3x T_{ij} = \frac{1}{2} \int d^3x \bar{T}_{00} x_i x_j = \frac{1}{2} \ddot{Q}_{ij}$$

# Energy loss in Minkowski

---




# Energy loss in Minkowski

## Energy and power

$$H_t = \frac{G}{8\pi} \int du \int d\Sigma_2 \ddot{\bar{Q}}_{ij}^{TT}(u) \ddot{\bar{Q}}_{TT}^{ij}(u)$$

$$P_t = \frac{G}{8\pi} \int d\Sigma_2 \ddot{\bar{Q}}_{ij}^{TT}(u) \ddot{\bar{Q}}_{TT}^{ij}(u)$$


where  $\dot{\bar{Q}}_{ij} = \mathcal{L}_t \bar{Q}_{ij}$ ,  
TT=Transverse & Traceless



## First ingredient: $h_{ab}$

---

$$g_{\mu\nu}(\epsilon) = a^2(\eta) (\mathring{g}_{\mu\nu} + \epsilon h_{\mu\nu} + \dots)$$



Need late time expansion instead of  $1/r$  expansion!

## First ingredient: $h_{ab}$

---

$$ds^2 = -dt^2 + e^{2Ht} (dx^2 + dy^2 + dz^2) \quad H^2 = \frac{\Lambda}{3}$$

## First ingredient: $h_{ab}$

$$ds^2 = -dt^2 + e^{2Ht} (dx^2 + dy^2 + dz^2) \quad H^2 = \frac{\Lambda}{3}$$

Gravitational field satisfies

$$\left( -\frac{\partial^2}{\partial t^2} + e^{-2Ht} D^2 - 3H \frac{\partial}{\partial t} \right) \bar{h}_{ij} = 16\pi G e^{-2Ht} T_{ij}$$

which has solutions in the late time regime [Vega, Ramirez, Sanchez; 1998]

$$\bar{h}_{ij} = \frac{4G}{r} \underbrace{\int_{-\infty}^{t_r} dt' e^{Ht'} \frac{\partial}{\partial t'} \int d^3x' T_{ij}(t', x')}_{\text{tail term}} + \mathcal{O}\left(\frac{D_o}{\ell_\Lambda} + \frac{D_o}{d(t)}\right)$$

## First ingredient: $h_{ab}$

$$ds^2 = -dt^2 + e^{2Ht} (dx^2 + dy^2 + dz^2) \quad H^2 = \frac{\Lambda}{3}$$

Gravitational field satisfies

$$\left( -\frac{\partial^2}{\partial t^2} + e^{-2Ht} D^2 - 3H \frac{\partial}{\partial t} \right) \bar{h}_{ij} = 16\pi G e^{-2Ht} T_{ij}$$

which has solutions in the late time regime [Vega, Ramirez, Sanchez; 1998]

$$\bar{h}_{ij} = \frac{4G}{r} \underbrace{\int d^3x' T_{ij}(t_r, x') - 4GH \int_{-\infty}^{t_r} dt' e^{Ht'} \frac{\partial}{\partial t'} \int d^3x' T_{ij}(t', x')}_{\text{tail term}} + \mathcal{O}\left(\frac{D_o}{\ell_\Lambda} + \frac{D_o}{d(t)}\right)$$

Assumptions

- ▶  $D(t) \leq D_o \ll \ell_\Lambda \equiv \frac{1}{H}$
- ▶  $D_o < d(t)$  (better for late times due to expansion)
- ▶  $\mathcal{L}_T T_{ab}$  is non-zero only for a finite time-interval

## Second ingredient: energy formula

---

Using the symplectic framework, the Hamiltonian associated with de Sitter time translation is

$$\begin{aligned} H_T &= -\frac{1}{2}\omega(h, \mathcal{L}_T h) \\ &= \frac{1}{16\pi GH} \int d^3\mathcal{I}^+ \mathcal{E}^{ij} \left( \mathcal{L}_T \bar{h}_{ij}^{TT} + 2H \bar{h}_{ij}^{TT} \right) \end{aligned}$$

## Third ingredient: quadrupole moment I

Definition

$$Q_{ij} := \underbrace{\int d^3V}_{a^3 dx dy dz} \underbrace{T_{\mu\nu} \partial_t^\mu \partial_t^\nu}_{\rho} \underbrace{(ax_i)(ax_j)}_{\text{physical distance}}$$

## Third ingredient: quadrupole moment I

Definition

$$Q_{ij} := \underbrace{\int d^3V}_{a^3 dx dy dz} \underbrace{\rho}_{T_{\mu\nu} \partial_t^\mu \partial_t^\nu} \underbrace{(ax_i)(ax_j)}_{\text{physical distance}}$$

Interesting feature of de Sitter:

$$\mathcal{L}_T T_{\mu\nu} = 0 \implies \mathcal{L}_T \rho = 0$$

$$\implies \mathcal{L}_T Q_{ij} = \int d^3V \cancel{\mathcal{L}_T \rho} (ax_i)(ax_j) - 2HQ_{ij} \xrightarrow{0} 0 \neq 0$$

## Third ingredient: quadrupole moment II

---

Conservation of stress-energy tensor  $\bar{\nabla}^\mu T_{\mu\nu} = 0$  is

$$\partial_t \rho - e^{-Ht} D^i T_{0i} + 3H(\rho + P) = 0$$

$$\partial_t T_{0i} - D^j T_{ij} + 3HT_{0i} = 0$$

$$\begin{aligned} \Rightarrow \int d^3x T_{ij} &= \frac{1}{2} \int d^3V \left( \ddot{\rho} + 3He^{-Ht}\dot{\rho} + 3H^2e^{-2Ht}\rho \right) (ax_i)(ax_j) \\ &\quad + \frac{1}{2} \int d^3V \left( He^{-Ht}\dot{P} + 3H^2e^{-2Ht}P \right) (ax_i)(ax_j) \end{aligned}$$

## Third ingredient: quadrupole moment II

---

Conservation of stress-energy tensor  $\bar{\nabla}^\mu T_{\mu\nu} = 0$  is

$$\partial_t \rho - e^{-Ht} D^i T_{0i} + 3H(\rho + P) = 0$$

$$\partial_t T_{0i} - D^j T_{ij} + 3HT_{0i} = 0$$

$$\begin{aligned}\implies \int d^3x T_{ij} &= \frac{1}{2} \int d^3V \left( \ddot{\rho} + 3He^{-Ht}\dot{\rho} + 3H^2e^{-2Ht}\rho \right) (ax_i)(ax_j) \\ &\quad + \frac{1}{2} \int d^3V \left( He^{-Ht}\dot{P} + 3H^2e^{-2Ht}P \right) (ax_i)(ax_j)\end{aligned}$$

## Third ingredient: quadrupole moment II

Conservation of stress-energy tensor  $\bar{\nabla}^\mu T_{\mu\nu} = 0$  is

$$\partial_t \rho - e^{-Ht} D^i T_{0i} + 3H(\rho + P) = 0$$


$$\partial_t T_{0i} - D^j T_{ij} + 3HT_{0i} = 0$$

$$\begin{aligned} \Rightarrow \int d^3x T_{ij} &= \frac{1}{2} \int d^3V \left( \ddot{\rho} + 3He^{-Ht}\dot{\rho} + 3H^2e^{-2Ht}\rho \right) (ax_i)(ax_j) \\ &\quad + \frac{1}{2} \int d^3V \left( He^{-Ht}\dot{P} + 3H^2e^{-2Ht}P \right) (ax_i)(ax_j) \end{aligned}$$

$$\int d^3x T_{ij} = \frac{a^{-1}}{2} \left( \ddot{Q}_{ij}^{(\rho)} + 2H\dot{Q}_{ij}^{(\rho)} + H\dot{Q}_{ij}^{(P)} + 2H^2Q_{ij}^{(P)} \right)$$

# Energy

$$H_T = \frac{G}{8\pi} \int d^3\mathcal{I}^+ (\mathcal{R}_{ab}\mathcal{R}_{cd}^{TT})_{ret} \mathring{q}^{ac} \mathring{q}^{bd}$$



$$\mathcal{R}_{ab} = \ddot{Q}_{ab}^{(\rho)} + 3H\ddot{Q}_{ab}^{(\rho)} + 2H^2\dot{Q}_{ab}^{(\rho)} + H\ddot{Q}_{ab}^{(P)} + 3H^2\dot{Q}_{ab}^{(P)} + 2H^3Q_{ab}^{(P)}$$


# Properties energy

---


$$H_T = \frac{G}{8\pi} \int d^3\mathcal{I}^+ (\mathcal{R}_{ab}\mathcal{R}_{cd}^{TT})_{ret} \mathring{q}^{ac} \mathring{q}^{bd}$$

- ▶ Features information about energy density *and* pressure
- ▶ Recover result Minkowski in limit  $H \rightarrow 0$
- ▶  $\mathcal{L}_T T_{ab} = 0 \implies \dot{Q}_{ab} \neq 0$  but  $H_T = 0$  (as it should!)
- ▶ Energy is positive

How does positivity arise?



How does positivity arise?



# Conclusion & future work

---

$$P_T = \frac{G}{8\pi} \int d\Sigma_2 (\mathcal{R}_{ab}\mathcal{R}_{cd}^{TT})_{ret} \mathring{q}^{ac}\mathring{q}^{bd}$$

- ▶ Important for conceptual understanding: one step closer to a complete framework of asymptotics with  $\Lambda > 0$
- ▶ Observational consequences: replace de Sitter background by a FLRW background with  $\Lambda > 0$
- ▶ Any suggestions?

## Conclusion & future work

---

$$P_T = \frac{G}{8\pi} \int d\Sigma_2 (\mathcal{R}_{ab}\mathcal{R}_{cd}^{TT})_{ret} \mathring{q}^{ac}\mathring{q}^{bd}$$

- ▶ Important for conceptual understanding: one step closer to a complete framework of asymptotics with  $\Lambda > 0$
- ▶ Observational consequences: replace de Sitter background by a FLRW background with  $\Lambda > 0$
- ▶ Any suggestions?

## Conclusion & future work

$$P_T = \frac{G}{8\pi} \int d\Sigma_2 (\mathcal{R}_{ab}\mathcal{R}_{cd}^{TT})_{ret} \mathring{q}^{ac}\mathring{q}^{bd}$$

- ▶ Important for conceptual understanding: one step closer to a complete framework of asymptotics with  $\Lambda > 0$
- ▶ Observational consequences: replace de Sitter background by a FLRW background with  $\Lambda > 0$
- ▶ Any suggestions?

# Conclusion & future work

---

$$P_T = \frac{G}{8\pi} \int d\Sigma_2 (\mathcal{R}_{ab}\mathcal{R}_{cd}^{TT})_{ret} \mathring{q}^{ac}\mathring{q}^{bd}$$

- ▶ Important for conceptual understanding: one step closer to a complete framework of asymptotics with  $\Lambda > 0$
- ▶ Observational consequences: replace de Sitter background by a FLRW background with  $\Lambda > 0$
- ▶ Any suggestions?

THANK YOU