

# A Toy Penrose Inequality and Its Proof

Emma Jakobsson  
Ingemar Bengtsson

GR21, New York 2016

# The Penrose Inequality

$$M \geq \sqrt{\frac{A}{16\pi}}$$

$M$  : mass


$A$  : area of surface

# The Penrose Inequality

$$M \geq \sqrt{\frac{A}{16\pi}}$$

$M$  : mass

$A$  : area of surface



Null shell version


Penrose (1973), Gibbons (1973), Tod (1985), Mars & Soria (2016)

# The Penrose Inequality

$$M \geq \sqrt{\frac{A}{16\pi}}$$

$M$  : mass

$A$  : area of surface



## Null shell version

A geometric property of surfaces  
in Minkowski space:

$$\oint \theta_+ dA \geq \sqrt{16\pi A}$$

$\theta_+$  : outer null expansion of  
surface

Penrose (1973), Gibbons (1973), Tod (1985), Mars & Soria (2016)


# The Penrose Inequality

$$M \geq \sqrt{\frac{A}{16\pi}}$$

$M$  : mass

$A$  : area of surface

Saturated by the event horizon of the Schwarzschild black hole.



## Null shell version

A geometric property of surfaces in Minkowski space:

$$\oint \theta_+ dA \geq \sqrt{16\pi A}$$

$\theta_+$  : outer null expansion of surface

Penrose (1973), Gibbons (1973), Tod (1985), Mars & Soria (2016)

# The Toy Version

The nonrotating BTZ black hole:

$$\ell^2 M = \left( \frac{L_{BTZ}}{2\pi} \right)^2$$

$M$  : mass of black hole

$\ell$  : length scale of  $\text{adS}_3$

$L_{BTZ}$  : length of event horizon

# The Toy Version

The nonrotating BTZ black hole:

$$\ell^2 M = \left( \frac{L_{BTZ}}{2\pi} \right)^2$$

$M$  : mass of black hole  
 $\ell$  : length scale of  $\text{adS}_3$   
 $L_{BTZ}$  : length of event horizon

Circular cross section of past light cone:

$$\frac{1}{2\pi} \oint \theta_+ dI = \ell^2 + \left( \frac{L}{2\pi} \right)^2$$

# The Toy Version

The nonrotating BTZ black hole:

$$\ell^2 M = \left( \frac{L_{BTZ}}{2\pi} \right)^2$$

$M$  : mass of black hole  
 $\ell$  : length scale of  $\text{adS}_3$   
 $L_{BTZ}$  : length of event horizon

Circular cross section of past light cone:


$$\frac{1}{2\pi} \oint \theta_+ dI = \ell^2 + \left( \frac{L}{2\pi} \right)^2$$

Toy version Penrose inequality in 2+1-dimensional anti-de Sitter space:

$$\frac{1}{2\pi} \oint \theta_+ dI \geq \ell^2 + \left( \frac{L}{2\pi} \right)^2$$




# The Proof



2+1-dimensional anti-de Sitter space as a stack of Poincaré disks:

$$ds^2 = -\ell^2 \left( \frac{1+\rho^2}{1-\rho^2} \right)^2 dt^2 + \frac{4\ell^2}{(1-\rho^2)^2} (d\rho^2 + \rho^2 d\phi^2)$$

# The Proof




## Null surface $\Sigma$

Characterized by its intersection with  $\mathcal{I}$ .

$$\gamma_{\mathcal{I}}: \quad \rho = 1, \quad \phi = \sigma, \quad t = T(\sigma)$$

# The Proof




## Null surface $\Sigma$

Characterized by its intersection with  $\mathcal{I}$ .

$$\gamma_{\mathcal{I}}: \quad \rho = 1, \quad \phi = \sigma, \quad t = T(\sigma)$$

Ruled by null geodesics parameterized by  $\tau$ .

# The Proof



## Null surface $\Sigma$

Characterized by its intersection with  $\mathcal{I}$ .


$$\gamma_{\mathcal{I}}: \quad \rho = 1, \quad \phi = \sigma, \quad t = T(\sigma)$$

Ruled by null geodesics parameterized by  $\tau$ .

## Cross section $\gamma$ of $\Sigma$

$$\gamma: \quad \tau = \tau(\sigma) \quad \text{on } \Sigma$$

# The Proof



## Null surface $\Sigma$

Characterized by its intersection with  $\mathcal{I}$ .

$$\gamma_{\mathcal{I}}: \quad \rho = 1, \quad \phi = \sigma, \quad t = T(\sigma)$$

Ruled by null geodesics parameterized by  $\tau$ .

## Cross section $\gamma$ of $\Sigma$

$$\gamma: \quad \tau = \tau(\sigma) \quad \text{on } \Sigma$$

$$\text{Length: } L = \ell \int_0^{2\pi} \cot[\tau(\sigma)] d\sigma$$

# The Proof

Proof.

Once  $\theta_+$  has been calculated we observe that

$$\frac{1}{2\pi} \oint_{\gamma} \theta_+ \, dl = \ell^2 \left( 1 + \int_0^{2\pi} \cot^2[\tau(\sigma)] \frac{d\sigma}{2\pi} + \int_0^{2\pi} f^2(\sigma) \frac{d\sigma}{2\pi} \right)$$

# The Proof

Proof.

Once  $\theta_+$  has been calculated we observe that

$$\begin{aligned}\frac{1}{2\pi} \oint_{\gamma} \theta_+ \, dl &= \ell^2 \left( 1 + \int_0^{2\pi} \cot^2[\tau(\sigma)] \frac{d\sigma}{2\pi} + \int_0^{2\pi} f^2(\sigma) \frac{d\sigma}{2\pi} \right) \\ &\geq \ell^2 + \left( \ell \int_0^{2\pi} \cot[\tau(\sigma)] \frac{d\sigma}{2\pi} \right)^2\end{aligned}$$

# The Proof

Proof.

Once  $\theta_+$  has been calculated we observe that

$$\begin{aligned}\frac{1}{2\pi} \oint_{\gamma} \theta_+ \, dl &= \ell^2 \left( 1 + \int_0^{2\pi} \cot^2[\tau(\sigma)] \frac{d\sigma}{2\pi} + \int_0^{2\pi} f^2(\sigma) \frac{d\sigma}{2\pi} \right) \\ &\geq \ell^2 + \left( \ell \int_0^{2\pi} \cot[\tau(\sigma)] \frac{d\sigma}{2\pi} \right)^2 \\ &= \ell^2 + \left( \frac{L}{2\pi} \right)^2\end{aligned}$$





Thank you for listening!