

Thinking outside the truncation: new hair for holographic superconductors

Gavin S. Hartnett

work with Jorge Santos and Harvey Reall, *work in progress*

Wednesday 13th July, 2016

GR21

Columbia University

Holographic Superconductors

Originally proposed by [Hartnoll, Herzog, Horowitz \(2008\)](#)

Consider 4D theory of Einstein-Maxwell-charged scalar

$$S = \int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F^2 - |\mathcal{D}\phi|^2 - V(\phi) \right), \quad \mathcal{D}_\mu = \nabla_\mu - iqA_\mu.$$

$$ds^2 = -f dt^2 + \frac{dr^2}{g} + r^2 (dx_1^2 + dx_2^2), \quad \phi = \phi(r). \quad (1)$$

For appropriate (q, V) , two classes of static, \mathbb{R}^2 -invariant solutions:

- 1 AdS-Reissner Nordstrom ($\phi = 0$)
- 2 charged AdS black hole with scalar hair ($\phi \neq 0$)

By considering more complicated matter content and geometries, this approach has been extended in many ways, by many authors

Some examples:

- Drude behaviour
- non-linear conductivity
- lattice effects
- cuprate-based superconductors
- momentum relaxation
- anisotropies/disorder

By considering more complicated matter content and geometries, this approach has been extended in many ways, by many authors

Some examples:

- Drude behaviour
- non-linear conductivity
- lattice effects
- cuprate-based superconductors
- momentum relaxation
- anisotropies/disorder

Bottom-up framework

It is interesting and important to consider these models in the context of precise AdS/CFT dualities

- bottom-up dualities are not fully specified or defined
- bottom-up models only retain subset of full 10/11D degrees of freedom

It is interesting and important to consider these models in the context of precise AdS/CFT dualities

- bottom-up dualities are not fully specified or defined
- bottom-up models only retain subset of full 10/11D degrees of freedom
- AdS/CMT dualities derivable from 10D string theory/11D M-theory
 - have the KK degrees of freedom
 - not freely specified – constrained by compactification

It is interesting and important to consider these models in the context of precise AdS/CFT dualities

- bottom-up dualities are not fully specified or defined
- bottom-up models only retain subset of full 10/11D degrees of freedom
- AdS/CMT dualities derivable from 10D string theory/11D M-theory
 - have the KK degrees of freedom
 - not freely specified – constrained by compactification
- Does requiring an AdS/CMT duality to be “top-down” lead to any important consequences/restrictions/effects?

Consider 11D SUGRA

$$2\kappa_{11}^2 S = \int d^{11}x \sqrt{-g} \left(R - \frac{1}{2} |G_4|^2 \right) + \frac{1}{6} \int C_3 \wedge G_4 \wedge G_4 ,$$

Top-down framework

Consider 11D SUGRA

$$2\kappa_{11}^2 S = \int d^{11}x \sqrt{-g} \left(R - \frac{1}{2} |G_4|^2 \right) + \frac{1}{6} \int C_3 \wedge G_4 \wedge G_4,$$

simple Freund-Rubin solution:

$$ds_{11}^2 = ds^2(AdS_4) + 4 \underbrace{[\eta \otimes \eta + ds^2(\mathbb{CP}^3)]}_{S^7}, \quad G_4 = -3\text{vol}(AdS_4).$$

Top-down framework

Consider 11D SUGRA

$$2\kappa_{11}^2 S = \int d^{11}x \sqrt{-g} \left(R - \frac{1}{2} |G_4|^2 \right) + \frac{1}{6} \int C_3 \wedge G_4 \wedge G_4 ,$$

simple Freund-Rubin solution:

$$ds_{11}^2 = ds^2(AdS_4) + 4 \underbrace{[\eta \otimes \eta + ds^2(\mathbb{CP}^3)]}_{S^7}, \quad G_4 = -3 \text{vol}(AdS_4) .$$

Top-down embedding of Holographic Superconductor [Gauntlett, Sonner, Wiseman \(2009\)](#)

$$ds_{11}^2 = e^{-4U} ds_4^2 + 4e^{2U} ds^2(\mathbb{CP}^3) + 4e^{-4U} (\eta + A_1) \otimes (\eta + A_1)$$

$$G_4 = -3e^{-12U} (1 - |\chi|^2) \text{vol}_4 + \frac{3i}{4} e^{-12U} \star_4 (\chi^* D\chi - \chi D\chi^*) \wedge (\eta + A_1) \\ + F_2 \wedge J + \sqrt{3} \left[\chi (\eta + A_1) \wedge \Omega - \frac{i}{4} D\chi \wedge \Omega + cc \right]$$

$$F = dA_1, \quad J = dA/2, \quad e^{6U} = 1 - \frac{3}{4} |\chi|^2$$

Top-down framework

Resulting non-linear 4D effective action:

$$S = \int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F^2 - h(\chi) |D\chi|^2 - V(\chi) \right)$$

Example of what's called a consistent truncation:

- only subset of all KK modes are retained
- Remarkable that this exists at all!

Expanded around $\chi = 0$,

$$S = \int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F^2 - |D\tilde{\chi}|^2 + 2|\tilde{\chi}|^2 \right)$$

$\tilde{\chi}$ has mass $M^2 = -2$ ($\Delta = 1, 2$) and $q = 4$

Top-down framework

Resulting non-linear 4D effective action:

$$S = \int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F^2 - h(\chi) |D\chi|^2 - V(\chi) \right)$$

Example of what's called a consistent truncation:

- only subset of all KK modes are retained
- Remarkable that this exists at all!

Expanded around $\chi = 0$,

$$S = \int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F^2 - |D\tilde{\chi}|^2 + 2|\tilde{\chi}|^2 \right)$$

$\tilde{\chi}$ has mass $M^2 = -2$ ($\Delta = 1, 2$) and $q = 4$

$\tilde{\chi}$ is a **pseudo-scalar**: $\Delta = 2$ required by supersymmetry
Breitenlohner, Freedman (1982)

Top-down framework

Resulting non-linear 4D effective action:

$$S = \int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F^2 - h(\chi) |D\chi|^2 - V(\chi) \right)$$

Example of what's called a consistent truncation:

- only subset of all KK modes are retained
- Remarkable that this exists at all!

Expanded around $\chi = 0$,

$$S = \int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F^2 - |D\tilde{\chi}|^2 + 2|\tilde{\chi}|^2 \right)$$

$\tilde{\chi}$ has mass $M^2 = -2$ ($\Delta = 1, 2$) and $q = 4$

$\tilde{\chi}$ is a **pseudo-scalar**: $\Delta = 2$ required by supersymmetry
Breitenlohner, Freedman (1982)



\Rightarrow unstable to developing scalar hair!

Superconducting Instabilities

near 2nd order phase transition, non-linearities negligible:

$$(\mathcal{D}^2 - M^2)\tilde{\chi} = 0$$

- $\tilde{\chi} = \exp(-i\omega t)\varphi(r)$
- $T > T_c$: only phase is Reissner Nordstrom (stable quasi-normal mode)
- $T = T_c$: linearized zero mode
- $T < T_c$: hairy BH phase (linear instability)





Superconducting Instabilities

near 2nd order phase transition, non-linearities negligible:

$$(\mathcal{D}^2 - M^2)\tilde{\chi} = 0$$



- $\tilde{\chi} = \exp(-i\omega t)\varphi(r)$
- $T > T_c$: only phase is Reissner Nordstrom (stable quasi-normal mode)
- $T = T_c$: linearized zero mode
- $T < T_c$: hairy BH phase (linear instability)



smaller Δ , larger $q \Rightarrow$ stronger instability (higher T_c)

Superconducting Instabilities



Δ	charge q	T_c/μ	found by
1	4	0.803688	
1	2	0.347821	
2	4	0.0831536	Gauntlett, Sonner, Wiseman (2009)
2	2	0.00707295	



smaller Δ , larger $q \Rightarrow$ stronger instability (higher T_c)

Superconducting Instabilities

Δ	charge q	T_c/μ	found by
1	4	0.803688	
1	2	0.347821	Donos, Gauntlett (2011)
2	4	0.0831536	Gauntlett, Sonner, Wiseman (2009)
2	2	0.00707295	



smaller Δ , larger $q \Rightarrow$ stronger instability (higher T_c)

- Are there any other instabilities?

General Perturbations of 11D Embedding

- Are there any other instabilities?
- we considered perturbations around 11D AdS-RN $\times S^7$ background

$$\delta R_{AB} - \frac{1}{6} \delta G_{(A|CDE} G_{|B)}{}^{CDE} + \frac{1}{144} h_{AB} G^2 + \frac{1}{4} G_{(A|ECD} G_{B)P}{}^{CD} h^{EP} \\ + \frac{g_{AB}}{144} \left(2G \cdot \delta G - 4G_{CDEF} G_G{}^{DEF} h^{CG} \right) = 0,$$

$$d \star \delta G_4 + d(\delta \star) G_4 - G_4 \wedge \delta G_4 = 0.$$

- in general won't fit in consistent truncation

General Perturbations of 11D Embedding

- Are there any other instabilities?
- we considered perturbations around 11D AdS-RN $\times S^7$ background

$$\delta R_{AB} - \frac{1}{6} \delta G_{(A|CDE} G_{|B)}{}^{CDE} + \frac{1}{144} h_{AB} G^2 + \frac{1}{4} G_{(A|ECD} G_{B)P}{}^{CD} h^{EP} \\ + \frac{g_{AB}}{144} \left(2G \cdot \delta G - 4G_{CDEF} G_G{}^{DEF} h^{CG} \right) = 0,$$

$$d \star \delta G_4 + d(\delta \star) G_4 - G_4 \wedge \delta G_4 = 0.$$

- in general won't fit in consistent truncation
- need to separate out the \mathbb{CP}^3 dependence: use charged \mathbb{CP}^3 harmonics (previously used in perturbations of Myers-Perry BHs)
- charged: $(\mathcal{L}_\eta - im)\delta g_{AB} = 0$
- from 4D perspective perturbation has charge $q = m$

General Perturbations of 11D Embedding

- Among many different \mathbb{CP}^3 harmonics, found one remarkably simple perturbation
- Out of δg_{AB} , δG_4 and their derivatives, constructed gauge invariant scalar Ψ obeying $(\mathcal{D}^2 - M^2) \Psi = 0$, with $M^2 = -2$ and $q = 4$

General Perturbations of 11D Embedding

- Among many different \mathbb{CP}^3 harmonics, found one remarkably simple perturbation
- Out of δg_{AB} , δG_4 and their derivatives, constructed gauge invariant scalar Ψ obeying $(\mathcal{D}^2 - M^2) \Psi = 0$, with $M^2 = -2$ and $q = 4$
- Ψ is a **scalar**, (rather than pseudo-scalar), so

General Perturbations of 11D Embedding

- Among many different \mathbb{CP}^3 harmonics, found one remarkably simple perturbation
- Out of δg_{AB} , δG_4 and their derivatives, constructed gauge invariant scalar Ψ obeying $(\mathcal{D}^2 - M^2) \Psi = 0$, with $M^2 = -2$ and $q = 4$
- Ψ is a **scalar**, (rather than pseudo-scalar), so
- $\Delta = (\textcolor{red}{1}, 2)$

General Perturbations of 11D Embedding

- Among many different \mathbb{CP}^3 harmonics, found one remarkably simple perturbation
- Out of δg_{AB} , δG_4 and their derivatives, constructed gauge invariant scalar Ψ obeying $(\mathcal{D}^2 - M^2)\Psi = 0$, with $M^2 = -2$ and $q = 4$
- Ψ is a **scalar**, (rather than pseudo-scalar), so
- $\Delta = (1, 2)$

Δ	charge q	T_c/μ	found by
1	4	0.803688	GSH, Reall, Santos
1	2	0.347821	Donos, Gauntlett (2011)
2	4	0.0831536	Gauntlett, Sonner, Wiseman (2009)
2	2	0.00707295	

Dominant instability/first mode to go unstable!!

- previous studies of this embedding constructed non-linear hairy solutions based off of sub-dominant instabilities

Implications & Further Directions

- previous studies of this embedding constructed non-linear hairy solutions based off of sub-dominant instabilities
 - these are sub-leading saddles, not relevant in thermodynamic limit
 - need to backreact Ψ & construct non-linear solution!

Implications & Further Directions

- previous studies of this embedding constructed non-linear hairy solutions based off of sub-dominant instabilities
 - these are sub-leading saddles, not relevant in thermodynamic limit
 - need to backreact Ψ & construct non-linear solution!
- seems unlikely that Ψ fits into a consistent truncation, in which case:

Implications & Further Directions

- previous studies of this embedding constructed non-linear hairy solutions based off of sub-dominant instabilities
 - these are sub-leading saddles, not relevant in thermodynamic limit
 - need to backreact Ψ & construct non-linear solution!
- seems unlikely that Ψ fits into a consistent truncation, in which case:
 - non-linear phase necessarily involves infinite set of KK modes
 - seems unusual from a condensed matter perspective \rightarrow implications for AdS/CMT?
 - familiar from attempts to model d-wave superconductivity holographically, [Hartnett, Horowitz \(2013\)](#), [Kim, Taylor \(2013\)](#)

Implications & Further Directions

- previous studies of this embedding constructed non-linear hairy solutions based off of sub-dominant instabilities
 - these are sub-leading saddles, not relevant in thermodynamic limit
 - need to backreact Ψ & construct non-linear solution!
- seems unlikely that Ψ fits into a consistent truncation, in which case:
 - non-linear phase necessarily involves infinite set of KK modes
 - seems unusual from a condensed matter perspective \rightarrow implications for AdS/CMT?
 - familiar from attempts to model d-wave superconductivity holographically, [Hartnett, Horowitz \(2013\)](#), [Kim, Taylor \(2013\)](#)
 - somewhat unusual: “lumpy” black holes do not often dominate grand canonical ensemble (c.f. [Santos/Way/Dias](#) talks)

- Application to landscape of superconducting membranes ([Denef, Hartnoll \(2009\)](#)): we specifically looked at the case $KE_6 = \mathbb{CP}^3$, how universal is this scalar Ψ (depends on spectrum of charged harmonics on KE_6)

Thank You!