

# A lower bound for the mass of multiple charged rotating black holes

Gilbert Weinstein  
Ariel University, Israel

GR21  
Columbia University, New York  
July 2016

# The Penrose heuristic argument

The standard picture of gravitational collapse: Any asymptotically flat initial data set evolving under the Einstein equations eventually settles to a number of Kerr-Newman solutions scattering away from each other.

# The Penrose heuristic argument

The standard picture of gravitational collapse: Any asymptotically flat initial data set evolving under the Einstein equations eventually settles to a number of Kerr-Newman solutions scattering away from each other.

Only three parameters per each of these black holes survive the collapse: the total mass  $m_i$ , angular momentum  $J_i$ , and charge  $q_i$ .

# The Penrose heuristic argument

The standard picture of gravitational collapse: Any asymptotically flat initial data set evolving under the Einstein equations eventually settles to a number of Kerr-Newman solutions scattering away from each other.

Only three parameters per each of these black holes survive the collapse: the total mass  $m_i$ , angular momentum  $J_i$ , and charge  $q_i$ . For a Kerr-Newman black hole we have:

$$m_i^2 = \frac{A_i}{16\pi} + \frac{q_i^2}{2} + \frac{\pi(q_i^4 + 4\mathcal{J}_i^2)}{A_i}$$

or minimizing over  $A_i$ :

$$m_i^2 \geq \frac{q_i^2 + \sqrt{q_i^4 + 4\mathcal{J}_i^2}}{2},$$

with equality iff the Kerr-Newman is *extreme*.

# Heuristic arguments (cont.)

Let  $m_0$ ,  $\mathcal{J}_0$ ,  $q_0$  denote the mass, angular momentum, and total charge of an initial state.

## Heuristic arguments (cont.)

Let  $m_0$ ,  $\mathcal{J}_0$ ,  $q_0$  denote the mass, angular momentum, and total charge of an initial state. Under appropriate hypotheses

$$\mathcal{J}_0 = \mathcal{J} = \sum \mathcal{J}_i, \quad q_0 = q = \sum q_i,$$

## Heuristic arguments (cont.)

Let  $m_0$ ,  $\mathcal{J}_0$ ,  $q_0$  denote the mass, angular momentum, and total charge of an initial state. Under appropriate hypotheses  $\mathcal{J}_0 = \mathcal{J} = \sum \mathcal{J}_i$ ,  $q_0 = q = \sum q_i$ , and  $m_0 \geq m = \sum m_i$ .

# Heuristic arguments (cont.)

Let  $m_0$ ,  $\mathcal{J}_0$ ,  $q_0$  denote the mass, angular momentum, and total charge of an initial state. Under appropriate hypotheses

$\mathcal{J}_0 = \mathcal{J} = \sum \mathcal{J}_i$ ,  $q_0 = q = \sum q_i$ , and  $m_0 \geq m = \sum m_i$ .

$$\sqrt{2}m = \sqrt{2} \sum m_i \geq \sum \sqrt{q_i^2 + \sqrt{q_i^4 + 4\mathcal{J}_i^2}} \geq \sqrt{q^2 + \sqrt{q^4 + 4\mathcal{J}^2}}.$$



# Heuristic arguments (cont.)

Let  $m_0$ ,  $\mathcal{J}_0$ ,  $q_0$  denote the mass, angular momentum, and total charge of an initial state. Under appropriate hypotheses

$\mathcal{J}_0 = \mathcal{J} = \sum \mathcal{J}_i$ ,  $q_0 = q = \sum q_i$ , and  $m_0 \geq m = \sum m_i$ .

$$\sqrt{2}m = \sqrt{2} \sum m_i \geq \sum \sqrt{q_i^2 + \sqrt{q_i^4 + 4\mathcal{J}_i^2}} \geq \sqrt{q^2 + \sqrt{q^4 + 4\mathcal{J}^2}}.$$

## Lemma

Let  $a_i, b_i \in \mathbb{R}$  and let  $a = \sum a_i$ ,  $b = \sum b_i$ . Then

$$(a^4 + b^2)^{1/4} \leq \sum (a_i^4 + b_i^2)^{1/4}.$$

# Heuristic arguments (cont.)

Let  $m_0$ ,  $\mathcal{J}_0$ ,  $q_0$  denote the mass, angular momentum, and total charge of an initial state. Under appropriate hypotheses

$\mathcal{J}_0 = \mathcal{J} = \sum \mathcal{J}_i$ ,  $q_0 = q = \sum q_i$ , and  $m_0 \geq m = \sum m_i$ .

$$\sqrt{2}m = \sqrt{2} \sum m_i \geq \sum \sqrt{q_i^2 + \sqrt{q_i^4 + 4\mathcal{J}_i^2}} \geq \sqrt{q^2 + \sqrt{q^4 + 4\mathcal{J}^2}}.$$

## Lemma

Let  $a_i, b_i \in \mathbb{R}$  and let  $a = \sum a_i$ ,  $b = \sum b_i$ . Then

$$(a^4 + b^2)^{1/4} \leq \sum (a_i^4 + b_i^2)^{1/4}.$$

## Lemma

Let  $a_i, b_i \in \mathbb{R}$  and let  $a = \sum a_i$ ,  $b = \sum b_i$ . Then

$$\sqrt{a^2 + \sqrt{a^4 + b^2}} \leq \sum \sqrt{a_i^2 + \sqrt{a_i^4 + b_i^2}}.$$

# $m, q, J$ inequality for multiple black holes

## Theorem (Khuri-W)

*Let  $(M, g, k, E, B)$  be a smooth, simply connected, axially symmetric, maximal initial data set satisfying  $\mu_{EM} \geq 0$  and  $g(J_{EM}, \eta) = 0$ , and with  $N + 1$  ends, one designated AF and the others either AF or AC. Then*

$$m \geq \mathcal{F}(\mathcal{J}_1, \dots, \mathcal{J}_N, q_1^e, \dots, q_N^e, q_1^b, \dots, q_N^b). \quad (1)$$

# $m, q, J$ inequality for multiple black holes

## Theorem (Khuri-W)

*Let  $(M, g, k, E, B)$  be a smooth, simply connected, axially symmetric, maximal initial data set satisfying  $\mu_{EM} \geq 0$  and  $g(J_{EM}, \eta) = 0$ , and with  $N + 1$  ends, one designated AF and the others either AF or AC. Then*

$$m \geq \mathcal{F}(\mathcal{J}_1, \dots, \mathcal{J}_N, q_1^e, \dots, q_N^e, q_1^b, \dots, q_N^b). \quad (1)$$

The function  $\mathcal{F}$  is the reduced energy of a harmonic map.

# $m, q, J$ inequality for multiple black holes

## Theorem (Khuri-W)

*Let  $(M, g, k, E, B)$  be a smooth, simply connected, axially symmetric, maximal initial data set satisfying  $\mu_{EM} \geq 0$  and  $g(J_{EM}, \eta) = 0$ , and with  $N + 1$  ends, one designated AF and the others either AF or AC. Then*

$$m \geq \mathcal{F}(\mathcal{J}_1, \dots, \mathcal{J}_N, q_1^e, \dots, q_N^e, q_1^b, \dots, q_N^b). \quad (1)$$

The function  $\mathcal{F}$  is the reduced energy of a harmonic map.

Work in progress:  $\mathcal{F} = \sqrt{\left(q^2 + \sqrt{q^4 + 4\mathcal{J}^2}\right)}/2$ .

# $m, q, J$ inequality for multiple black holes

## Theorem (Khuri-W)

*Let  $(M, g, k, E, B)$  be a smooth, simply connected, axially symmetric, maximal initial data set satisfying  $\mu_{EM} \geq 0$  and  $g(J_{EM}, \eta) = 0$ , and with  $N + 1$  ends, one designated AF and the others either AF or AC. Then*

$$m \geq \mathcal{F}(\mathcal{J}_1, \dots, \mathcal{J}_N, q_1^e, \dots, q_N^e, q_1^b, \dots, q_N^b). \quad (1)$$

The function  $\mathcal{F}$  is the reduced energy of a harmonic map.

Work in progress:  $\mathcal{F} = \sqrt{\left(q^2 + \sqrt{q^4 + 4\mathcal{J}^2}\right)}/2$ .

## Conjecture

*Equality cannot be achieved if  $N > 1$  unless all charges are of the same sign and the angular momenta vanish. In this case, the initial data set is isometric to the canonical slice of a MP spacetime.*

# Previous Work

- ▶ S. Dain, *Proof of the angular momentum-mass inequality for axisymmetric black hole*, JDG, **79** (2008), 33-67.

# Previous Work

- ▶ S. Dain, *Proof of the angular momentum-mass inequality for axisymmetric black hole*, JDG, **79** (2008), 33-67.
- ▶ S. Dain, *Geometric inequalities for axially symmetric black holes*, CQG, **29** (2012), no. 7, 073001.



# Previous Work

- ▶ S. Dain, *Proof of the angular momentum-mass inequality for axisymmetric black hole*, JDG, **79** (2008), 33-67.
- ▶ S. Dain, *Geometric inequalities for axially symmetric black holes*, CQG, **29** (2012), no. 7, 073001.
- ▶ P. Chruściel, *Mass and angular-momentum inequalities for axi-symmetric initial data sets. I. Positivity of Mass*, Ann. Phys., **323** (2008), 2566-2590.

# Previous Work

- ▶ S. Dain, *Proof of the angular momentum-mass inequality for axisymmetric black hole*, JDG, **79** (2008), 33-67.
- ▶ S. Dain, *Geometric inequalities for axially symmetric black holes*, CQG, **29** (2012), no. 7, 073001.
- ▶ P. Chruściel, *Mass and angular-momentum inequalities for axi-symmetric initial data sets. I. Positivity of Mass*, Ann. Phys., **323** (2008), 2566-2590.
- ▶ P. Chruściel, Y. Li, and G. Weinstein, *Mass and angular-momentum inequalities for axi-symmetric initial data sets. II. Angular Momentum*, Ann. Phys., **323** (2008), 2591-2613.

# Previous Work

- ▶ S. Dain, *Proof of the angular momentum-mass inequality for axisymmetric black hole*, JDG, **79** (2008), 33-67.
- ▶ S. Dain, *Geometric inequalities for axially symmetric black holes*, CQG, **29** (2012), no. 7, 073001.
- ▶ P. Chruściel, *Mass and angular-momentum inequalities for axi-symmetric initial data sets. I. Positivity of Mass*, Ann. Phys., **323** (2008), 2566-2590.
- ▶ P. Chruściel, Y. Li, and G. Weinstein, *Mass and angular-momentum inequalities for axi-symmetric initial data sets. II. Angular Momentum*, Ann. Phys., **323** (2008), 2591-2613.
- ▶ R. Schoen, and X. Zhou, *Convexity of reduced energy and mass angular momentum inequalities*, Ann. Henri Poincaré, **14** (2013), 1747-1773.

# Previous Work

- ▶ S. Dain, *Proof of the angular momentum-mass inequality for axisymmetric black hole*, JDG, **79** (2008), 33-67.
- ▶ S. Dain, *Geometric inequalities for axially symmetric black holes*, CQG, **29** (2012), no. 7, 073001.
- ▶ P. Chruściel, *Mass and angular-momentum inequalities for axi-symmetric initial data sets. I. Positivity of Mass*, Ann. Phys., **323** (2008), 2566-2590.
- ▶ P. Chruściel, Y. Li, and G. Weinstein, *Mass and angular-momentum inequalities for axi-symmetric initial data sets. II. Angular Momentum*, Ann. Phys., **323** (2008), 2591-2613.
- ▶ R. Schoen, and X. Zhou, *Convexity of reduced energy and mass angular momentum inequalities*, Ann. Henri Poincaré, **14** (2013), 1747-1773.
- ▶ P. Chruściel, and J. Costa, *Mass, angular-momentum and charge inequalities for axisymmetric initial data*, CQG, **26** (2009), no. 23, 235013.

# Previous Work

- ▶ S. Dain, *Proof of the angular momentum-mass inequality for axisymmetric black hole*, JDG, **79** (2008), 33-67.
- ▶ S. Dain, *Geometric inequalities for axially symmetric black holes*, CQG, **29** (2012), no. 7, 073001.
- ▶ P. Chruściel, *Mass and angular-momentum inequalities for axi-symmetric initial data sets. I. Positivity of Mass*, Ann. Phys., **323** (2008), 2566-2590.
- ▶ P. Chruściel, Y. Li, and G. Weinstein, *Mass and angular-momentum inequalities for axi-symmetric initial data sets. II. Angular Momentum*, Ann. Phys., **323** (2008), 2591-2613.
- ▶ R. Schoen, and X. Zhou, *Convexity of reduced energy and mass angular momentum inequalities*, Ann. Henri Poincaré, **14** (2013), 1747-1773.
- ▶ P. Chruściel, and J. Costa, *Mass, angular-momentum and charge inequalities for axisymmetric initial data*, CQG, **26** (2009), no. 23, 235013.
- ▶ J. Costa, *Proof of a Dain inequality with charge*, J. Phys. A, **43** (2010), no. 28, 285202.

# Sketch of proof

Under the hypotheses, one may define EM potentials  $\chi$ ,  $\psi$ , and a twist potential  $\nu$ , so that

$$m \geq \mathcal{M}(U, \nu, \chi, \psi) = \frac{1}{8\pi} \int_{\mathbb{R}^3} \left( |\nabla U|^2 + \frac{e^{4U}}{\rho^4} |\nabla \nu + \chi \nabla \psi - \psi \nabla \chi|^2 + \frac{e^{2U}}{\rho^2} (|\nabla \chi|^2 + |\nabla \psi|^2) \right) dx$$

# Sketch of proof

Under the hypotheses, one may define EM potentials  $\chi$ ,  $\psi$ , and a twist potential  $v$ , so that

$$m \geq \mathcal{M}(U, v, \chi, \psi) = \frac{1}{8\pi} \int_{\mathbb{R}^3} \left( |\nabla U|^2 + \frac{e^{4U}}{\rho^4} |\nabla v + \chi \nabla \psi - \psi \nabla \chi|^2 + \frac{e^{2U}}{\rho^2} (|\nabla \chi|^2 + |\nabla \psi|^2) \right) dx$$

If we define  $u = U - \log \rho$ , then

$$\mathcal{H}_\Omega(u, v, \chi, \psi) = \mathcal{M}_\Omega(U, v, \chi, \psi) + \int_\Omega \frac{|\nabla \rho|^2}{\rho^2} - \int_\Omega \nabla U \cdot \nabla \log \rho,$$

# Sketch of proof

Under the hypotheses, one may define EM potentials  $\chi$ ,  $\psi$ , and a twist potential  $v$ , so that

$$m \geq \mathcal{M}(U, v, \chi, \psi) = \frac{1}{8\pi} \int_{\mathbb{R}^3} \left( |\nabla U|^2 + \frac{e^{4U}}{\rho^4} |\nabla v + \chi \nabla \psi - \psi \nabla \chi|^2 + \frac{e^{2U}}{\rho^2} (|\nabla \chi|^2 + |\nabla \psi|^2) \right) dx$$

If we define  $u = U - \log \rho$ , then

$$\mathcal{H}_\Omega(u, v, \chi, \psi) = \mathcal{M}_\Omega(U, v, \chi, \psi) + \int_\Omega \frac{|\nabla \rho|^2}{\rho^2} - \int_\Omega \nabla U \cdot \nabla \log \rho,$$

$\mathcal{H}$  is the harmonic map energy for  $\Phi = (u, v, \psi, \chi): \mathbb{R}^3 \rightarrow \mathbb{H}_{\mathbb{C}}^2$ .



# Sketch of proof

Under the hypotheses, one may define EM potentials  $\chi$ ,  $\psi$ , and a twist potential  $v$ , so that

$$m \geq \mathcal{M}(U, v, \chi, \psi) = \frac{1}{8\pi} \int_{\mathbb{R}^3} \left( |\nabla U|^2 + \frac{e^{4U}}{\rho^4} |\nabla v + \chi \nabla \psi - \psi \nabla \chi|^2 + \frac{e^{2U}}{\rho^2} (|\nabla \chi|^2 + |\nabla \psi|^2) \right) dx$$

If we define  $u = U - \log \rho$ , then

$$\mathcal{H}_\Omega(u, v, \chi, \psi) = \mathcal{M}_\Omega(U, v, \chi, \psi) + \int_\Omega \frac{|\nabla \rho|^2}{\rho^2} - \int_\Omega \nabla U \cdot \nabla \log \rho,$$

$\mathcal{H}$  is the harmonic map energy for  $\Phi = (u, v, \psi, \chi): \mathbb{R}^3 \rightarrow \mathbb{H}_{\mathbb{C}}^2$ .

Since  $\log \rho$  is harmonic, the EL equations for  $\mathcal{H}_\Omega(u, v, \chi, \psi)$ , and  $\mathcal{M}_\Omega(u, v, \chi, \psi)$  are the same.

# Sketch of proof

The proof consists in showing

# Sketch of proof

The proof consists in showing

- ▶ There is a critical point of  $\mathcal{M}(U, \nu, \chi, \psi)$  with the appropriate boundary conditions and asymptotics. (Harmonic map with prescribed singularities)

# Sketch of proof

The proof consists in showing

- ▶ There is a critical point of  $\mathcal{M}(U, \nu, \chi, \psi)$  with the appropriate boundary conditions and asymptotics. (Harmonic map with prescribed singularities)
- ▶ The critical point is a minimizer of  $\mathcal{M}$ . (Convexity under geodesic deformation)

# Sketch of proof

The proof consists in showing

- ▶ There is a critical point of  $\mathcal{M}(U, \nu, \chi, \psi)$  with the appropriate boundary conditions and asymptotics. (Harmonic map with prescribed singularities)
- ▶ The critical point is a minimizer of  $\mathcal{M}$ . (Convexity under geodesic deformation)

The function  $\mathcal{F}$  is then the reduced energy  $\mathcal{M}$  of that minimizer.

# Sketch of proof

The proof consists in showing

- ▶ There is a critical point of  $\mathcal{M}(U, v, \chi, \psi)$  with the appropriate boundary conditions and asymptotics. (Harmonic map with prescribed singularities)
- ▶ The critical point is a minimizer of  $\mathcal{M}$ . (Convexity under geodesic deformation)

The function  $\mathcal{F}$  is then the reduced energy  $\mathcal{M}$  of that minimizer.

The boundary conditions are:

$$\mathcal{J}_n = \frac{1}{4}(v|_{I_n} - v|_{I_{n-1}}),$$

# Sketch of proof

The proof consists in showing

- ▶ There is a critical point of  $\mathcal{M}(U, v, \chi, \psi)$  with the appropriate boundary conditions and asymptotics. (Harmonic map with prescribed singularities)
- ▶ The critical point is a minimizer of  $\mathcal{M}$ . (Convexity under geodesic deformation)

The function  $\mathcal{F}$  is then the reduced energy  $\mathcal{M}$  of that minimizer.

The boundary conditions are:

$$\mathcal{J}_n = \frac{1}{4}(v|_{I_n} - v|_{I_{n-1}}),$$

$$q_n^e = \frac{1}{2}(\chi|_{I_n} - \chi|_{I_{n-1}}), \quad q_n^b = \frac{1}{2}(\psi|_{I_n} - \psi|_{I_{n-1}}),$$

# Sketch of proof

The proof consists in showing

- ▶ There is a critical point of  $\mathcal{M}(U, v, \chi, \psi)$  with the appropriate boundary conditions and asymptotics. (Harmonic map with prescribed singularities)
- ▶ The critical point is a minimizer of  $\mathcal{M}$ . (Convexity under geodesic deformation)

The function  $\mathcal{F}$  is then the reduced energy  $\mathcal{M}$  of that minimizer.

The boundary conditions are:

$$\mathcal{J}_n = \frac{1}{4}(v|_{I_n} - v|_{I_{n-1}}),$$

$$q_n^e = \frac{1}{2}(\chi|_{I_n} - \chi|_{I_{n-1}}), \quad q_n^b = \frac{1}{2}(\psi|_{I_n} - \psi|_{I_{n-1}}),$$

where  $I_n$  is the interval of the  $z$ -axis  $\Gamma$  from  $p_n$  to  $p_{n+1}$ ,  $p_0 = -\infty$  and  $p_{N+1} = \infty$ .



# Existence of the harmonic map

The tension of a map  $\Phi = (u, v, \chi, \psi)$  is:

$$\tau^u = \Delta u - 2e^{4u}|\nabla v + \chi\nabla\psi - \psi\nabla\chi|^2 + e^{2u}(|\nabla\chi|^2 + |\nabla\psi|^2),$$

$$\begin{aligned}\tau^v = & \Delta v + 2\nabla u \cdot \nabla v + 2(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot \nabla u \\ & - 2e^{2u}(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot (\chi\nabla\chi + \psi\nabla\psi),\end{aligned}$$

$$\tau^\chi = \Delta\chi + 2\nabla u \cdot \nabla\chi - 2e^{2u}(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot \nabla\psi,$$

$$\tau^\psi = \Delta\psi + 2\nabla u \cdot \nabla\psi + 2e^{2u}(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot \nabla\chi.$$

# Existence of the harmonic map

The tension of a map  $\Phi = (u, v, \chi, \psi)$  is:

$$\tau^u = \Delta u - 2e^{4u}|\nabla v + \chi\nabla\psi - \psi\nabla\chi|^2 + e^{2u}(|\nabla\chi|^2 + |\nabla\psi|^2),$$

$$\begin{aligned}\tau^v = & \Delta v + 2\nabla u \cdot \nabla v + 2(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot \nabla u \\ & - 2e^{2u}(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot (\chi\nabla\chi + \psi\nabla\psi),\end{aligned}$$

$$\tau^\chi = \Delta\chi + 2\nabla u \cdot \nabla\chi - 2e^{2u}(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot \nabla\psi,$$

$$\tau^\psi = \Delta\psi + 2\nabla u \cdot \nabla\psi + 2e^{2u}(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot \nabla\chi.$$

Thus  $\tau(\Phi) = 0$  iff  $\Phi$  is harmonic.

# Existence of the harmonic map

The tension of a map  $\Phi = (u, v, \chi, \psi)$  is:

$$\tau^u = \Delta u - 2e^{4u}|\nabla v + \chi\nabla\psi - \psi\nabla\chi|^2 + e^{2u}(|\nabla\chi|^2 + |\nabla\psi|^2),$$

$$\begin{aligned}\tau^v = & \Delta v + 2\nabla u \cdot \nabla v + 2(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot \nabla u \\ & - 2e^{2u}(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot (\chi\nabla\chi + \psi\nabla\psi),\end{aligned}$$

$$\tau^\chi = \Delta\chi + 2\nabla u \cdot \nabla\chi - 2e^{2u}(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot \nabla\psi,$$


$$\tau^\psi = \Delta\psi + 2\nabla u \cdot \nabla\psi + 2e^{2u}(\nabla v + \chi\nabla\psi - \psi\nabla\chi) \cdot \nabla\chi.$$

Thus  $\tau(\Phi) = 0$  iff  $\Phi$  is harmonic.

## Lemma

*There is a map  $\tilde{\Phi}_0$ , such that its reduced energy is finite, its tension  $\tau(\tilde{\Phi}_0)$  has support inside a bounded set, and  $\tau(\tilde{\Phi}_0)$  is pointwise bounded. Moreover, the values of  $(\tilde{v}_0, \tilde{\chi}_0, \tilde{\psi}_0)$  agree with those of the given data  $(v, \chi, \psi)$  on each component of*

*$\Gamma' = \Gamma \setminus \{p_1, \dots, p_N\}$ , and  $\tilde{U}_0 = \tilde{u}_0 + \log \rho = \log r_n + o_{I-4}(r_n^{1/2})$  near each puncture  $p_n$ .*



# Existence of the harmonic map

We say that two maps  $\Phi, \Psi: \mathbb{R}^3 \setminus \Gamma \rightarrow \mathbb{H}_{\mathbb{C}}^2$  are *asymptotic* if  $\text{dist}_{\mathbb{H}_{\mathbb{C}}^2}(\Phi, \Psi)$  is bounded.

# Existence of the harmonic map

We say that two maps  $\Phi, \Psi: \mathbb{R}^3 \setminus \Gamma \rightarrow \mathbb{H}_{\mathbb{C}}^2$  are *asymptotic* if  $\text{dist}_{\mathbb{H}_{\mathbb{C}}^2}(\Phi, \Psi)$  is bounded.

## Corollary

*For any set of punctures  $p_n$  on the axis  $\Gamma$  and prescribed constants  $v_0|_{I_n}, \chi_0|_{I_n}, \psi_0|_{I_n}, n = 1, \dots, N$ , there exists a corresponding unique harmonic map  $\tilde{\Psi}_0 = (u_0, v_0, \chi_0, \psi_0)$  which is asymptotic to  $\tilde{\Phi}_0$ , and satisfies*

$$U_0 = u_0 + \log \rho = \log r_n + O(1), \quad v_0, \chi_0, \psi_0 = O(1), \quad \text{as } r_n \rightarrow 0.$$

# Existence of the harmonic map

We say that two maps  $\Phi, \Psi: \mathbb{R}^3 \setminus \Gamma \rightarrow \mathbb{H}_{\mathbb{C}}^2$  are *asymptotic* if  $\text{dist}_{\mathbb{H}_{\mathbb{C}}^2}(\Phi, \Psi)$  is bounded.

## Corollary

*For any set of punctures  $p_n$  on the axis  $\Gamma$  and prescribed constants  $v_0|_{I_n}, \chi_0|_{I_n}, \psi_0|_{I_n}, n = 1, \dots, N$ , there exists a corresponding unique harmonic map  $\tilde{\Psi}_0 = (u_0, v_0, \chi_0, \psi_0)$  which is asymptotic to  $\tilde{\Phi}_0$ , and satisfies*

$$U_0 = u_0 + \log \rho = \log r_n + O(1), \quad v_0, \chi_0, \psi_0 = O(1), \quad \text{as } r_n \rightarrow 0.$$

*Furthermore, on  $\mathbb{R}^3 \setminus \Gamma$ , the reduced energy density  $\mathcal{E}'(\tilde{\Psi}_0) \leq c\rho^{-2}$ .*

# Existence of the harmonic map

We say that two maps  $\Phi, \Psi: \mathbb{R}^3 \setminus \Gamma \rightarrow \mathbb{H}_{\mathbb{C}}^2$  are *asymptotic* if  $\text{dist}_{\mathbb{H}_{\mathbb{C}}^2}(\Phi, \Psi)$  is bounded.

## Corollary

*For any set of punctures  $p_n$  on the axis  $\Gamma$  and prescribed constants  $v_0|_{I_n}, \chi_0|_{I_n}, \psi_0|_{I_n}, n = 1, \dots, N$ , there exists a corresponding unique harmonic map  $\tilde{\Psi}_0 = (u_0, v_0, \chi_0, \psi_0)$  which is asymptotic to  $\tilde{\Phi}_0$ , and satisfies*

$$U_0 = u_0 + \log \rho = \log r_n + O(1), \quad v_0, \chi_0, \psi_0 = O(1), \quad \text{as } r_n \rightarrow 0.$$

*Furthermore, on  $\mathbb{R}^3 \setminus \Gamma$ , the reduced energy density  $\mathcal{E}'(\tilde{\Psi}_0) \leq c\rho^{-2}$ .*

The proof of the corollary uses the convexity of  $\text{dist}_{\mathbb{H}_{\mathbb{C}}^2}$ .



# Existence of the harmonic map

We say that two maps  $\Phi, \Psi: \mathbb{R}^3 \setminus \Gamma \rightarrow \mathbb{H}_{\mathbb{C}}^2$  are *asymptotic* if  $\text{dist}_{\mathbb{H}_{\mathbb{C}}^2}(\Phi, \Psi)$  is bounded.

## Corollary

*For any set of punctures  $p_n$  on the axis  $\Gamma$  and prescribed constants  $v_0|_{I_n}, \chi_0|_{I_n}, \psi_0|_{I_n}, n = 1, \dots, N$ , there exists a corresponding unique harmonic map  $\tilde{\Psi}_0 = (u_0, v_0, \chi_0, \psi_0)$  which is asymptotic to  $\tilde{\Phi}_0$ , and satisfies*

$$U_0 = u_0 + \log \rho = \log r_n + O(1), \quad v_0, \chi_0, \psi_0 = O(1), \quad \text{as } r_n \rightarrow 0.$$

*Furthermore, on  $\mathbb{R}^3 \setminus \Gamma$ , the reduced energy density  $\mathcal{E}'(\tilde{\Psi}_0) \leq c\rho^{-2}$ .*

The proof of the corollary uses the convexity of  $\text{dist}_{\mathbb{H}_{\mathbb{C}}^2}$ .

- G. Weinstein, *Harmonic maps with prescribed singularities into Hadamard manifolds*, Mathematical Research Letters, **3** (1996), no 6, 835-844.

# Thank you!