

Causality theory for $C^{1,1}$ metrics

James Vickers

joint work with

M. Kunzinger

R. Steinbauer

M. Stojković

**21st International Conference
on General Relativity and Gravitation**

C^2 - Causality Theory

C^2 - Causality Theory

- Majority of standard references on causality theory [1] assume that the metric is smooth, or at least sufficiently differentiable for all proofs to work.
- Most other treatments [2] are more explicit about the differentiability of the metric and assume the metric is C^2 .
- Only a few [3] consider lower regularity metrics.

C^2 - Causality Theory

- Majority of standard references on causality theory [1] assume that the metric is smooth, or at least sufficiently differentiable for all proofs to work.
- Most other treatments [2] are more explicit about the differentiability of the metric and assume the metric is C^2 .
- Only a few [3] consider lower regularity metrics.
- For C^2 metrics the inverse function theorem shows that locally the causal structure is as in Minkowski space, in the sense that

$$\exp_p : \tilde{U} \rightarrow U \quad \text{is a } C^1\text{-diffeomorphism}$$

and

$$I^+(p) \cap U = \exp_p \left(I^+(0) \cap \tilde{U} \right)$$

For metrics below C^2 this result is problematic.

Why consider metrics of regularity lower than C^2 ?

Why consider metrics of regularity lower than C^2 ?

- (1) **PDE Theory:** (Classical result) [Choquet-Bruhat] There exist solutions of the Einstein vacuum equations with initial data $(h, k) \in H_{\text{loc}}^s \times H_{\text{loc}}^{s-1}$ with $s > 5/2$ for which g is not C^2 . Recent results using bilinear estimates allow solutions of even lower regularity c.f. the L^2 -curvature conjecture. [Klainermann et al]

Why consider metrics of regularity lower than C^2 ?

- (1) **PDE Theory:** (Classical result) [Choquet-Bruhat] There exist solutions of the Einstein vacuum equations with initial data $(h, k) \in H_{\text{loc}}^s \times H_{\text{loc}}^{s-1}$ with $s > 5/2$ for which g is not C^2 . Recent results using bilinear estimates allow solutions of even lower regularity c.f. the L^2 -curvature conjecture. [Klainermann et al]
- (2) **Physical models:** Want to be able to include situations where there is a shock in the matter or a jump in the energy-momentum tensor across an interface, such as the boundary of a star.

jump in density \rightarrow jump in curvature \rightarrow metric not C^2

Why consider metrics of regularity lower than C^2 ?

- (1) **PDE Theory:** (Classical result) [Choquet-Bruhat] There exist solutions of the Einstein vacuum equations with initial data $(h, k) \in H_{\text{loc}}^s \times H_{\text{loc}}^{s-1}$ with $s > 5/2$ for which g is not C^2 . Recent results using bilinear estimates allow solutions of even lower regularity c.f. the L^2 -curvature conjecture. [Klainermann et al]
- (2) **Physical models:** Want to be able to include situations where there is a shock in the matter or a jump in the energy-momentum tensor across an interface, such as the boundary of a star.

jump in density \rightarrow jump in curvature \rightarrow metric not C^2

- (3) **Singularity Theorems:** Failure to be C^2 need not be very singular (see below and next talk). So we want to be able to prove singularity theorems in lower regularity than C^2 . Need causality theory in lower regularity than C^2 .

Why $C^{1,1}$ metrics?

$C^{1,1}$ metrics are continuously differentiable with Lipschitz first derivatives. By Rademacher's Theorem $\partial_i g_{jk}$ is differentiable almost everywhere and as a result $C^{1,1}$ is often written as C^{2-} .

Why $C^{1,1}$ metrics?

$C^{1,1}$ metrics are continuously differentiable with Lipschitz first derivatives. By Rademacher's Theorem $\partial_i g_{jk}$ is differentiable almost everywhere and as a result $C^{1,1}$ is often written as C^{2-} .

$C^{1,1}$ metrics have the following properties:

- Exist unique local solutions to the geodesic equation
- $\exp_p : \tilde{U} \rightarrow U$ is Lipschitz
- The curvature is defined almost everywhere and is bounded.

Why $C^{1,1}$ metrics?

$C^{1,1}$ metrics are continuously differentiable with Lipschitz first derivatives. By Rademacher's Theorem $\partial_i g_{jk}$ is differentiable almost everywhere and as a result $C^{1,1}$ is often written as C^{2-} .

$C^{1,1}$ metrics have the following properties:

- Exist unique local solutions to the geodesic equation
- $\exp_p : \tilde{U} \rightarrow U$ is Lipschitz
- The curvature is defined almost everywhere and is bounded.

Problems below $C^{1,1}$:

- Below $C^{1,1}$ geodesic convexity may no longer hold [Hartman & Wintner]
- Below $C^{0,1}$ may not be able to deform a causal curve that is not everywhere null into a timelike curve. (Pushup lemma fails) [Chrusciel & Grant].
- Below $C^{0,1}$ exist “bubbling metrics” with light-cones having non-empty interior [Chrusciel & Grant].

Totally convex neighbourhoods for $C^{1,1}$ metrics

Key results: [Kunzinger, Steinbauer, Stojković], [Minguzzi]

Totally convex neighbourhoods for $C^{1,1}$ metrics

Key results: [Kunzinger, Steinbauer, Stojković], [Minguzzi]

Theorem: Let M be a smooth manifold with a $C^{1,1}$ -pseudo-Riemannian metric g and let $p \in M$. Then there exist open neighbourhoods \tilde{U} of $0 \in T_p M$ and U of p in M such that

$$\exp_p : \tilde{U} \rightarrow U$$

is a *bi-Lipschitz homeomorphism*.

Remark: It follows from Rademacher's theorem that both \exp_p and \exp_p^{-1} are differentiable almost everywhere.

Totally convex neighbourhoods for $C^{1,1}$ metrics

Key results: [Kunzinger, Steinbauer, Stojković], [Minguzzi]

Theorem: Let M be a smooth manifold with a $C^{1,1}$ -pseudo-Riemannian metric g and let $p \in M$. Then there exist open neighbourhoods \tilde{U} of $0 \in T_p M$ and U of p in M such that

$$\exp_p : \tilde{U} \rightarrow U$$

is a *bi-Lipschitz homeomorphism*.

Remark: It follows from Rademacher's theorem that both \exp_p and \exp_p^{-1} are differentiable almost everywhere.

Theorem: Let M be a smooth manifold with a $C^{1,1}$ -pseudo-Riemannian metric g . Then each point $p \in M$ possesses a basis of totally normal neighbourhoods.

Remark: A totally normal set is geodesically convex.

Two complementary approaches to proving causality results for $C^{1,1}$ metrics

Two complementary approaches to proving causality results for $C^{1,1}$ metrics

- (1) [KSSV] Follow method of Chrusciel & Grant and approximate g by a family of smooth metrics $g_\varepsilon \rightarrow g$ while controlling the causal structure.

Two complementary approaches to proving causality results for $C^{1,1}$ metrics

- (1) [KSSV] Follow method of Chrusciel & Grant and approximate g by a family of smooth metrics $g_\varepsilon \rightarrow g$ while controlling the causal structure.
- (2) [Minguzzi] Look at geodesic equation for g and use Picard-Lindelöf iteration and an inverse function theorem for Lipschitz maps.

Two complementary approaches to proving causality results for $C^{1,1}$ metrics

- (1) [KSSV] Follow method of Chrusciel & Grant and approximate g by a family of smooth metrics $g_\varepsilon \rightarrow g$ while controlling the causal structure.
 - ▶ Can be applied to a wide class of metrics (not just $C^{1,1}$).
 - ▶ Can make use of results from smooth causality theory.
 - ▶ Less precise information.
 - ▶ Some results are valid only almost everywhere.
- (2) [Minguzzi] Look at geodesic equation for g and use Picard-Lindelöf iteration and an inverse function theorem for Lipschitz maps.
 - ▶ Gives more precise information e.g. $D_0 \exp_p = \text{id}_{T_p M}$
 - ▶ \exp is a bi-Lipschitz map in a neighbourhood of the zero section of TM
 - ▶ Not applicable below $C^{1,1}$.
 - ▶ Cannot make use of results from smooth theory.


Two complementary approaches to proving causality results for $C^{1,1}$ metrics

- (1) [KSSV] Follow method of Chrusciel & Grant and approximate g by a family of smooth metrics $g_\varepsilon \rightarrow g$ while controlling the causal structure.
 - ▶ Can be applied to a wide class of metrics (not just $C^{1,1}$).
 - ▶ Can make use of results from smooth causality theory.
 - ▶ Less precise information.
 - ▶ Some results are valid only almost everywhere.
- (2) [Minguzzi] Look at geodesic equation for g and use Picard-Lindelöf iteration and an inverse function theorem for Lipschitz maps.
 - ▶ Gives more precise information e.g. $D_0 \exp_p = \text{id}_{T_p M}$
 - ▶ \exp is a bi-Lipschitz map in a neighbourhood of the zero section of TM
 - ▶ Not applicable below $C^{1,1}$.
 - ▶ Cannot make use of results from smooth theory.

Both needed to prove singularity theorems in low regularity.

Chruściel-Grant regularisation of the metric


Sandwich null cones of g between null cones of two families of smooth metrics so that $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$.



Chruściel-Grant regularisation of the metric


Sandwich null cones of g between null cones of two families of smooth metrics so that $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$.

- Locally, $g_\varepsilon := g * \rho_\varepsilon$, glued by partition of unity subordinate to $U_i \subseteq M$.



Chruściel-Grant regularisation of the metric


Sandwich null cones of g between null cones of two families of smooth metrics so that $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$.



- Locally, $g_\varepsilon := g * \rho_\varepsilon$, glued by partition of unity subordinate to $U_i \Subset M$.
- Same way: construct time-like 1-form ω s.t. $|\omega(X)| \geq c_i > 0$ for all g -causal vector fields X with $\|X\|_h = 1$.

Chruściel-Grant regularisation of the metric


Sandwich null cones of g between null cones of two families of smooth metrics so that $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$.



- Locally, $g_\varepsilon := g * \rho_\varepsilon$, glued by partition of unity subordinate to $U_i \Subset M$.
- Same way: construct time-like 1-form ω s.t. $|\omega(X)| \geq c_i > 0$ for all g -causal vector fields X with $\|X\|_h = 1$.
- $\check{g}_{\eta,\lambda} := g_\eta + \lambda \omega \otimes \omega$

Chruściel-Grant regularisation of the metric

Sandwich null cones of g between null cones of two families of smooth metrics so that $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$.




- Locally, $g_\varepsilon := g * \rho_\varepsilon$, glued by partition of unity subordinate to $U_i \subseteq M$.
- Same way: construct time-like 1-form ω s.t. $|\omega(X)| \geq c_i > 0$ for all g -causal vector fields X with $\|X\|_h = 1$.
- $\check{g}_{\eta,\lambda} := g_\eta + \lambda\omega \otimes \omega$
- Adapt $\lambda = \lambda(\varepsilon)$ and $\eta = \eta(\varepsilon)$ locally s.t. for ε small

$$g(X, X) \leq 0 \ \& \ \|X\|_h = 1 \Rightarrow g_{\eta,\lambda}(X, X) < 0$$

Chruściel-Grant regularisation of the metric

Sandwich null cones of g between null cones of two families of smooth metrics so that $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$.




- Locally, $g_\varepsilon := g * \rho_\varepsilon$, glued by partition of unity subordinate to $U_i \Subset M$.
- Same way: construct time-like 1-form ω s.t. $|\omega(X)| \geq c_i > 0$ for all g -causal vector fields X with $\|X\|_h = 1$.
- $\check{g}_{\eta,\lambda} := g_\eta + \lambda\omega \otimes \omega$
- Adapt $\lambda = \lambda(\varepsilon)$ and $\eta = \eta(\varepsilon)$ locally s.t. for ε small

$$g(X, X) \leq 0 \ \& \ \|X\|_h = 1 \Rightarrow g_{\eta,\lambda}(X, X) < 0$$

- Glue w.r.t. x and ε to obtain Lorentzian metric $\check{g}_\varepsilon \prec g$.

Chruściel-Grant regularisation of the metric

Sandwich null cones of g between null cones of two families of smooth metrics so that $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$.



- Locally, $g_\varepsilon := g * \rho_\varepsilon$, glued by partition of unity subordinate to $U_i \Subset M$.
- Same way: construct time-like 1-form ω s.t. $|\omega(X)| \geq c_i > 0$ for all g -causal vector fields X with $\|X\|_h = 1$.
- $\check{g}_{\eta,\lambda} := g_\eta + \lambda\omega \otimes \omega$
- Adapt $\lambda = \lambda(\varepsilon)$ and $\eta = \eta(\varepsilon)$ locally s.t. for ε small

$$g(X, X) \leq 0 \ \& \ \|X\|_h = 1 \Rightarrow g_{\eta,\lambda}(X, X) < 0$$

- Glue w.r.t. x and ε to obtain Lorentzian metric $\check{g}_\varepsilon \prec g$.
- Similarly obtain \hat{g}_ε with $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$.

Properties of approximating metrics

- Given a Lorentzian metric of some prescribed regularity (e.g., Sobolev, Hölder, etc.), the convergence of the inner and outer regularisations \check{g}_ε and \hat{g}_ε to g is as good locally as that of regularisation by convolution.
- If g is a metric of general pseudo-Riemannian signature, then we may produce regularisations \check{g}_ε that are pseudo-Riemannian metrics on all of M of the same signature as g
- Every point has a basis of normal neighbourhoods U with $\exp_p : \tilde{U} \rightarrow U$, such that for ε sufficiently small, all $\exp_p^{g_\varepsilon}$ are diffeomorphisms with domain \tilde{U} . Moreover, the inverse maps $(\exp_p^{g_\varepsilon})^{-1}$ also are defined on a common neighbourhood of p for ε small, and converge locally uniformly to \exp_p^{-1} .

$C^{1,1}$ causality results

The following is the main local causality result for $C^{1,1}$ metrics:

Theorem: [KSSV] Let g be a $C^{1,1}$ -Lorentzian metric, and let $p \in M$. Then p has a basis of normal neighbourhoods U , $\exp_p : \tilde{U} \rightarrow U$ a bi-Lipschitz homeomorphism, such that:

$$I^+(p, U) = \exp_p(I^+(0) \cap \tilde{U})$$

$$J^+(p, U) = \exp_p(J^+(0) \cap \tilde{U})$$

$$\partial I^+(p, U) = \partial J^+(p, U) = \exp_p(\partial I^+(0) \cap \tilde{U})$$

Proof: ▶ Proof of Theorem

Remark: We follow [Chrusciel] in that we base our approach to causality theory on locally Lipschitz curves. This definition differs from that of [Minguzzi] (and others) where the corresponding curves are required to be C^1 . However we have the following Corollary to the above result:

Corollary: Let $U \subseteq M$ be open, $p \in U$. Then the sets $I^+(p, U)$, $J^+(p, U)$ remain unchanged if Lipschitz curves are replaced by piecewise C^1 curves, or in fact by broken geodesics.

References

[1] R. Penrose, *Techniques of Differential Topology in Relativity*, (1972)

Other causality references which assume a smooth metric are: O’Neil; Beem & Ehrlich; Minguzzi & Sanchez; and Kriele.

[2] P. T. Chruściel, *Elements of Causality Theory*, (2011)

Other causality references which only assume a C^2 metric are: Hawking & Ellis; and Senovilla.

[3] P. T. Chruściel, J. D. E. Grant,

On Lorentzian causality with continuous metrics,

Classical Quantum Gravity 29, 145001, 32 pp. (2012)

A. García-Parrado, J. M. M. Senovilla,

Causal structures and causal boundaries,

Classical Quantum Gravity 22, R1-R84, (2005)

Other causality references which consider lower regularity are: Carter; Clarke; Sorkin & Woolgar; Dowker, Garcia & Surya.

Y. Choquet Bruhat,

Theoreme d’Existence pour certains systemes d’equations aux derivees partielles nonlineaires,

Acta Math. 88, 141-225 (1952).

S. Klainerman, I. Rodnianski, J. Szeftel,

Overview of the proof of the bounded L^2 curvature conjecture,

arXiv:1204.1772 [gr-qc], (2012).

P. Hartman, A. Wintner,

On the problems of geodesics in the small,

Amer. J. Math. 73, 132–148 (1951).

M. Kunzinger, R. Steinbauer, M. Stojković.

The exponential map of a $C^{1,1}$ -metric, Differential Geom. Appl. 34, 14–24. (2014)

E. Minguzzi

Convex neighborhoods for Lipschitz connections and sprays,

Monatsh. Math. 177, no. 4, 569–625 (2015)

M. Kunzinger, R. Steinbauer, M. Stojković, J. A. Vickers.

A regularisation approach to causality theory for $C^{1,1}$ -Lorentzian metrics,

Gen. Relativity Gravitation 46, 1738 (2014)

Proof of $J^+(p, U) = \exp_p(J^+(0) \cap \tilde{U})$

Proof of $J^+(p, U) = \exp_p(J^+(0) \cap \tilde{U})$

1. We first show that $J^+(p, U) \supseteq \exp_p(J^+(0) \cap \tilde{U})$.

Let $v \in \tilde{U}$ and let $\alpha(t) = \exp_p(tv)$. Set $\alpha_\varepsilon(t) := \exp_p^{\hat{g}_\varepsilon}(tv)$.

Then $\alpha_\varepsilon \rightarrow \alpha$ in C^1 .

Hence applying the smooth Gauss lemma for each ε it follows that:

$$g(\alpha'(t), \alpha'(t)) = \lim_{\varepsilon \rightarrow 0} \hat{g}_\varepsilon(\alpha'_\varepsilon(t), \alpha'_\varepsilon(t)) = \lim_{\varepsilon \rightarrow 0} (\hat{g}_\varepsilon)_p(v, v) = g_p(v, v).$$

So $v \in \tilde{U} \cap J^+(0) \Rightarrow \exp_p(v) \in J^+(p, U)$.

Proof of $J^+(p, U) = \exp_p(J^+(0) \cap \tilde{U})$

1. We first show that $J^+(p, U) \supseteq \exp_p(J^+(0) \cap \tilde{U})$.

Let $v \in \tilde{U}$ and let $\alpha(t) = \exp_p(tv)$. Set $\alpha_\varepsilon(t) := \exp_p^{\hat{g}_\varepsilon}(tv)$.

Then $\alpha_\varepsilon \rightarrow \alpha$ in C^1 .

Hence applying the smooth Gauss lemma for each ε it follows that:

$$g(\alpha'(t), \alpha'(t)) = \lim_{\varepsilon \rightarrow 0} \hat{g}_\varepsilon(\alpha'_\varepsilon(t), \alpha'_\varepsilon(t)) = \lim_{\varepsilon \rightarrow 0} (\hat{g}_\varepsilon)_p(v, v) = g_p(v, v).$$

So $v \in \tilde{U} \cap J^+(0) \Rightarrow \exp_p(v) \in J^+(p, U)$.

2. We now show that $J^+(p, U) \subseteq \exp_p(J^+(0) \cap \tilde{U})$.

Let $\alpha : [0, 1] \rightarrow U$ be a FD causal curve in U from p

Let $\beta := (\exp_p)^{-1} \circ \alpha$ be the corresponding curve in $\tilde{U} \subset T_p M$.

Note that α is TL wrt \hat{g}_ε and set $\beta_\varepsilon := (\exp_{\hat{p}}^{\hat{g}_\varepsilon})^{-1} \circ \alpha$.

Then by *smooth causality* $\beta_\varepsilon([0, 1]) \subseteq I_{\hat{g}_\varepsilon(p)}^+(0)$ for all $\varepsilon < \varepsilon_0$.

Let $\tilde{Q}(v) = g_p(v, v)$ and $\tilde{Q}_\varepsilon(v) = (\hat{g}_\varepsilon)_p(v, v)$ be quadratic forms on $T_p M$.

Then $\beta_\varepsilon \rightarrow \beta$ uniformly, and $\tilde{Q}_\varepsilon \rightarrow \tilde{Q}$ locally uniformly.

So $\tilde{Q}(\beta(t)) = \lim_{\varepsilon \rightarrow 0} \tilde{Q}_\varepsilon(\beta_\varepsilon(t)) \leq 0$ Hence $\beta((0, 1]) \subseteq J^+(0) \cap \tilde{U}$.