

Impact of higher order modes in aligned spin searches for Binary Black Holes

PHYSICAL REVIEW D **93**, 084019 (2016)

Juan Calderón Bustillo


Georgia Institute of Technology

in collaboration with S.Husa, A.M. Sintes and M. Pürrer



GR21 Conference, Columbia University, New York, July 2016

Higher modes of BBH radiation





$$h_p(\Xi; r, \theta, \varphi; t) = h_+ - ih_\times = \frac{1}{d_L} \sum_{\ell \geq 2} \sum_{m=-\ell}^{m=\ell} Y_{\ell,m}^{-2}(\theta, \varphi) h_{\ell,m}(\Xi; t)$$

Modes: source info.

Harmonics: orientation info.


Searches do only include the $(\ell, |m|)=(2,2)$ modes



[JCB + (2016)]

Higher Modes are strong for.....

PN



NR (SXS)




FIGURE 2.5: Relative T1 (Top) and NR (Bottom) amplitude of the higher modes (3,3) and (4,4) relative to the dominant (2,2) mode as a function of the frequency for several non-spinning systems. Note how the higher the mass ratio q , the larger the contribution from higher order modes.





FIGURE 2.6: Relative T1 (Top) and NR (Bottom) amplitude of several higher modes relative to the dominant (2,2) mode as a function of the frequency for several $q=3$ spinning systems. Note that the influence of the spin is much lower than that due to the mass ratio.

[JCB + (2016)]


- Large mass ratio (q) BBH show stronger higher modes.
- Spin has a secondary effect.

Harmonics Impact (Orientation)




- (2,2) harmonic has its maximum when face-on and its minimum when edge on.
- Most higher harmonics behave in the opposite way.
- Edge-on systems have stronger higher mode content.

Total mass (M) and detector noise curve



**Higher modes important for high mass sources
(+ high mass ratio, + edge-on)**

Getting some intuition



Waveforms: Hybrid NR (SXS collaboration) + post Newtonian
Taylor T1

Hybridisation procedure: [JCB + (2015), arXiv:1501.00918]

Data analysis tools


Compute the fitting factor of hybrid PN+NR waveforms towards SEOBNRv1_ROM (only 2,2 mode)

$$\langle h|g \rangle = 4\Re \int_{f_0}^{\infty} \frac{\tilde{h}(f)\tilde{g}^*(f)}{S_n(f)} df$$

$$\mathcal{O} = \frac{\langle h|g \rangle}{\sqrt{\langle h|h \rangle \langle g|g \rangle}}$$

(2,2)-model: SEOBNRv1_ROM

[Pürrer + (2014)]



Average results weighting by optimal SNR

Maximum overlap with (2,2) model

M

q

% Event loss


$$R_i = \sum_j \left(\frac{\rho_{i,j} \mathcal{F}_{i,j}}{\rho_{i,j}} \right)^3$$

Parameter bias

$$\Xi_i^{\mathcal{B}} = \sum_j \Xi_{i,j}^{\mathcal{B}} \left(\frac{\rho_{i,j} \mathcal{F}_{i,j}}{\rho_{i,j}} \right)^3$$


$$\Delta \Xi_i = \Xi_{i,0} - \Xi_i^{\mathcal{B}}$$

Previous studies I




- Low match for edge-on orientations.
- These produce however the lowest SNR (for suitable low masses)


[L. Pekowsky + (2012)]



[Varma + (2014)]



Previous studies II



Reduction of **sensitivity** due to the larger number of templates.


Worth to include higher modes for certain part of the parameter space.

[Capano + (2013)]


- All considered the design Advanced LIGO curve ($f_0=10\text{Hz}$).
- Restricted to non-spinning targets and template banks
- We extend to aligned-spin searches and early Advanced LIGO ($f_0=30\text{Hz}$).

Warning: how to read sky-plots

- Sky around source projected onto the (x,y) orbital plane.
- **Center** corresponds to **face-on**.
- **Perimeter** corresponds to **edge-on**.
- The color code indicates the value of the measured magnitude.




Results: fitting factors vs. SNR



Results: fitting factors

- Fitting factors are lower the larger q and M are.
- Lower for edge-on cases.
- Lowest for iLIGO ($f=30\text{Hz}$ & non-spinning templates.)
- Larger for eaLIGO (spinning templates)
- Largest for aLIGO ($f=10\text{Hz.}$)



Results: averaged event loss





FIGURE 6.6: Top: Fractional volume loss in % for non-spinning $q = (4, 6, 8)$ systems in (dotted, dashed, solid). Bottom: same for $(q, \chi) = (3; 0, \pm 0.5)$ with the style code indicated in the corresponding caption. Note that since we used a non-spinning template bank for iLIGO, we did not consider spinning targets.

- 10% loss for $q > 4$ ($M > 100$ solar masses.) for AdvLIGO (note: $M = 100$ solar masses for non-spinning bank).
- 10% loss for $q > 4$ ($M > 100$ solar masses.) and $q > 6$ for all mass range for eaLIGO.
- Losses up to 25% for eaLIGO
- Very similar losses for spinning and non-spinning targets (spin is subdominant).


Results: Parameter Bias (total mass)

- Biases larger for edge-on systems and eaLIGO.
- Larger the larger q and M .
- Largest biases are towards lower total masses, as this increments the frequency of the bank waveform.
- When the mass is over-estimated, the mass ratio is underestimated and viceversa.



Results: Parameter Bias (spin)

- Biases larger for edge-on systems and eaLIGO.
- Larger the larger q and M .
- Largest biases are towards lower spins.
- Biases can get to -1 for non-spinning edge-on systems.



Results: averaged parameter bias

- Biases are larger the larger q and M are.
- Larger for eaLIGO ($f=30\text{Hz}$).
- Large spin biases for eaLIGO.
- Larger biases for negative spin systems.

[JCB + (2016)]





FIGURE 6.9: Top: M , M_c , and χ systematic bias for the $q = (3, 4, 6, 8)$ (from solid to dot-dashed) nS cases. We use the same color-detector code as in Fig.2. Bottom: Same for the $(q; \chi) = (3; 0, \pm 0.5)$. We use dashed (dotted) for - (+) spin and add $(q, \chi) = (1, -0.2)$ case in solid green for eaLIGO.

Conclusions & further work


- Higher modes have a large impact in systems with large mass ratio and total mass, specially if edge-on.
- Including a spin parameter in the template bank, reduces losses for the case of non-spinning targets (before applying signal based vetoes) at the cost of large spin bias.
- Losses of events (up to 26%) are larger for the case of early Advanced LIGO.
- Including higher modes is (in principle) worth for a larger portion of the parameter space for early advanced LIGO.
- Include precessing sources (in prep. with D. Shoemaker, P. Laguna K. Jani)
- Impact for chi-squared in gravitational wave search pipelines (in prep. with K.Jani)
- Build template bank including higher modes and run a search (addressing it...)

Precessing Targets I

- Precession dominates for low mass systems, and Design Advanced LIGO.
- Higher modes dominate for large total mass.




Precessing Targets II




Searches and chi-squared

Separation of injections and background via chi-squared

Injections do not contain
higher modes



Injections contain
higher modes



Injected signals: EOBNRv2 and EOBNRv2HM. $q=6$ $M=200$

Matched Filtering

Inner product



$$\langle h|g \rangle = 4\Re \int_{f_0}^{\infty} \frac{\tilde{h}(f)\tilde{g}^*(f)}{S_n(f)} df$$

SNR

$$\rho = \frac{\langle s|h \rangle}{\sqrt{\langle h|h \rangle}}$$

Overlap

$$\mathcal{O} = \frac{\langle h|g \rangle}{\sqrt{\langle h|h \rangle \langle g|g \rangle}}$$



Fitting Factor

Searches compare the incoming signal to a full bank B of templates h_i


Fitting Factor

$$FF = \text{Max}_i \mathcal{O}(h|h_i^B)$$

$$d \sim \rho(h|B) = FF \times \rho(h|h)$$

$$d \sim FF$$

$$V \sim FF^3$$



LVC+Virgo [2016]

Accessible volume proportional to FF^3

Note: There will be a bias on the recovered parameters

Results: comparison to statistical errors

