

# Parametrized theories: Making EM even “gaugier”

Juan Margalef Bentabol



Joint work with F. Barbero (CSIC) and Eduardo J.S. Villaseñor (UC3M)

21st International Conference on  
General Relativity and Gravitation



July 2016 - New York

## Motivation: parametrized

$(M \cong \Sigma \times I, g)$  being     $\Sigma$  3-manifold  
 $I = [t_0, t_1]$




## Motivation: parametrized

$(M \cong \Sigma \times I, g)$  being  $\Sigma$  3-manifold  
 $I = [t_0, t_1]$






# Motivation: parametrized

$(M \cong \Sigma \times I, g)$  being  $\Sigma$  3-manifold  
 $I = [t_0, t_1]$




# Motivation: parametrized

$(M \cong \Sigma \times I, g)$  being  $\Sigma$  3-manifold  
 $I = [t_0, t_1]$




# Motivation: parametrized

$(M \cong \Sigma \times I, g)$  being  $\Sigma$  3-manifold  
 $I = [t_0, t_1]$



# Motivation: parametrized

$(M \cong \Sigma \times I, g)$  being  $\Sigma$  3-manifold  
 $I = [t_0, t_1]$



$$Diff_{sp}^+(M) := \left\{ Z : M \rightarrow M \text{ diff.} / \begin{array}{l} TZ \cdot \partial_t \\ Z(\Sigma \times \{t\}) \end{array} \begin{array}{l} \text{future timelike} \\ \text{spacelike} \end{array} \right\}$$

# Our model

# Parametrized EM Field Action

$$S_{EM} : \Omega^1(M) \longrightarrow \mathbb{R}$$

$$S_{EM}[A] = \int_M (\mathrm{d}A) \wedge (\star_g \mathrm{d}A)$$

$$\mathrm{d} \star_g (\mathrm{d}A) = 0$$

# Parametrized EM Field Action

$$S_{EM} : \Omega^1(M) \longrightarrow \mathbb{R}$$

$$S_{EM}[A] = \int_M (\mathrm{d}A) \wedge (\star_g \mathrm{d}A)$$

$$\mathrm{d} \star_g (\mathrm{d}A) = 0$$

$$S_{EM}^P : \Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M) \longrightarrow \mathbb{R}$$

$$S_{EM}^P[A, Z] := \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A)$$

$$\mathrm{d} \star_{Z^*g} (\mathrm{d}A) = 0$$

# Parametrized EM Field Action

$$S_{EM} : \Omega^1(M) \longrightarrow \mathbb{R}$$

$$S_{EM}[A] = \int_M (\mathrm{d}A) \wedge (\star_g \mathrm{d}A)$$

$$\mathrm{d} \star_g (\mathrm{d}A) = 0$$

$$S_{EM}^P : \Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M) \longrightarrow \mathbb{R}$$

$$S_{EM}^P[A, Z] := \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A)$$

$$\mathrm{d} \star_{Z^*g} (\mathrm{d}A) = 0$$

## Properties

- If  $(A, Z)$  is a solution, then so is  $(Y^*A, Z \circ Y)$ .

# Parametrized EM Field Action

$$S_{EM} : \Omega^1(M) \longrightarrow \mathbb{R}$$

$$S_{EM}[A] = \int_M (\mathrm{d}A) \wedge (\star_g \mathrm{d}A)$$

$$\mathrm{d} \star_g (\mathrm{d}A) = 0$$

$$S_{EM}^P : \Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M) \longrightarrow \mathbb{R}$$

$$S_{EM}^P[A, Z] := \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A)$$

$$\mathrm{d} \star_{Z^*g} (\mathrm{d}A) = 0$$

## Properties

- If  $(A, Z)$  is a solution, then so is  $(Y^*A, Z \circ Y)$ .
- $S_{EM}^P[Y^*A, Z \circ Y] = S_{EM}^P[A, Z]$

# Hamiltonian Framework

# Hamiltonian Framework

$$S_{EM}^P = \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A)$$

$$M \cong \Sigma \times I$$

# Hamiltonian Framework

$$\left| \begin{array}{l} S_{EM}^P = \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A) \\ M \cong \Sigma \times I \end{array} \right. \xrightarrow{3+1} S_{EM}^P = \int_I \mathrm{d}t \int_\Sigma (something)_t$$

# Hamiltonian Framework

$$\left| \begin{array}{l} S_{EM}^P = \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A) \\ M \cong \Sigma \times I \end{array} \right. \xrightarrow{3+1} S_{EM}^P = \int_I \mathrm{d}t \int_\Sigma (something)_t$$

- Lagrangian  $L_{EM}^P : \mathcal{D} \subset TQ \rightarrow \mathbb{R}$

# Hamiltonian Framework

$$\left| \begin{array}{l} S_{EM}^P = \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A) \\ M \cong \Sigma \times I \end{array} \right. \xrightarrow{3+1} S_{EM}^P = \int_I \mathrm{d}t \int_\Sigma (something)_t$$

- Lagrangian  $L_{EM}^P : \mathcal{D} \subset TQ \rightarrow \mathbb{R}$
- Fiber derivative  $FL_{EM}^P : \mathcal{D} \subset TQ \rightarrow T^*Q$

# Hamiltonian Framework

$$\left| \begin{array}{l} S_{EM}^P = \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A) \\ M \cong \Sigma \times I \end{array} \right. \xrightarrow{3+1} S_{EM}^P = \int_I \mathrm{d}t \int_\Sigma (something)_t$$

- Lagrangian  $L_{EM}^P : \mathcal{D} \subset TQ \rightarrow \mathbb{R}$
- Fiber derivative  $FL_{EM}^P : \mathcal{D} \subset TQ \rightarrow T^*Q$
- Hamiltonian  $H_{EM}^P : FL_{EM}^P(\mathcal{D}) \rightarrow \mathbb{R}$

# Hamiltonian Framework

$$\left| \begin{array}{l} S_{EM}^P = \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A) \quad \xrightarrow{3+1} \quad S_{EM}^P = \int_I \mathrm{d}t \int_\Sigma (something)_t \\ M \cong \Sigma \times I \end{array} \right.$$

- Lagrangian  $L_{EM}^P : \mathcal{D} \subset TQ \rightarrow \mathbb{R}$
- Fiber derivative  $FL_{EM}^P : \mathcal{D} \subset TQ \rightarrow T^*Q$
- Hamiltonian  $H_{EM}^P : FL_{EM}^P(\mathcal{D}) \rightarrow \mathbb{R}$

## Important things to retain

- $L_{EM}^P$  homogeneous of degree 1, hence  $H_{EM}^P = 0$ .

# Hamiltonian Framework

$$\left| \begin{array}{l} S_{EM}^P = \int_M (\mathrm{d}A) \wedge (\star_{Z^*g} \mathrm{d}A) \quad \xrightarrow{3+1} \quad S_{EM}^P = \int_I \mathrm{d}t \int_\Sigma (\text{something})_t \\ M \cong \Sigma \times I \end{array} \right.$$

- Lagrangian  $L_{EM}^P : \mathcal{D} \subset TQ \rightarrow \mathbb{R}$
- Fiber derivative  $FL_{EM}^P : \mathcal{D} \subset TQ \rightarrow T^*Q$
- Hamiltonian  $H_{EM}^P : FL_{EM}^P(\mathcal{D}) \rightarrow \mathbb{R}$

## Important things to retain

- $L_{EM}^P$  homogeneous of degree 1, hence  $H_{EM}^P = 0$ .
- In fact, this is true for any parametrized theory.


# The geometric arena

# Configuration space

$$\Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M)$$

$$q^{(4)} : \Sigma \times I \rightarrow T^*(\Sigma \times I)$$

$$Z : \Sigma \times I \rightarrow \Sigma \times I$$




# Configuration space

$$\Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M)$$

$$q^{(4)} : \Sigma \times I \rightarrow T^*(\Sigma \times I)$$

$$Z : \Sigma \times I \rightarrow \Sigma \times I$$

$$X_t : \Sigma \hookrightarrow \Sigma \times I$$




# Configuration space

$$\Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M)$$

$$q^{(4)} : \Sigma \times I \rightarrow T^*(\Sigma \times I) \quad (q_\perp)_t : \Sigma \rightarrow \mathbb{R}$$

$$Z : \Sigma \times I \rightarrow \Sigma \times I$$

$$X_t : \Sigma \hookrightarrow \Sigma \times I$$



# Configuration space


$$\Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M)$$

$$q^{(4)} : \Sigma \times I \rightarrow T^*(\Sigma \times I)$$

$$\begin{aligned}(q_\perp)_t : \Sigma &\rightarrow \mathbb{R} \\ q_t : \Sigma &\rightarrow T^*\Sigma\end{aligned}$$

$$Z : \Sigma \times I \rightarrow \Sigma \times I$$

$$X_t : \Sigma \hookrightarrow \Sigma \times I$$



# Configuration space

$$\Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M)$$


$$Q_{EM} = C^\infty(\Sigma) \times \Omega^1(\Sigma) \times \text{Emb}(\Sigma, M)$$

$$q^{(4)} : \Sigma \times I \rightarrow T^*(\Sigma \times I)$$

$$\begin{aligned}(q_\perp)_t &: \Sigma \rightarrow \mathbb{R} \\ q_t &: \Sigma \rightarrow T^*\Sigma\end{aligned}$$

$$Z : \Sigma \times I \rightarrow \Sigma \times I$$

$$X_t : \Sigma \hookrightarrow \Sigma \times I$$



# Configuration space

$$\Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M)$$

$$Q_{EM} = C^\infty(\Sigma) \times \Omega^1(\Sigma) \times \text{Emb}(\Sigma, M)$$


$$q^{(4)} : \Sigma \times I \rightarrow T^*(\Sigma \times I)$$

$$(q_\perp)_t : \Sigma \rightarrow \mathbb{R}$$

$$q_t : \Sigma \rightarrow T^*\Sigma$$

$$Z : \Sigma \times I \rightarrow \Sigma \times I$$

$$X_t : \Sigma \hookrightarrow \Sigma \times I$$



# Configuration space

$$\Omega^1(M) \times \text{Diff}_{\text{sp}}^+(M)$$

$$Q_{EM} = C^\infty(\Sigma) \times \Omega^1(\Sigma) \times \text{Emb}_{\text{sp}}(\Sigma, M)$$


$$q^{(4)} : \Sigma \times I \rightarrow T^*(\Sigma \times I)$$

$$(q_\perp)_t : \Sigma \rightarrow \mathbb{R}$$

$$q_t : \Sigma \rightarrow T^*\Sigma$$

$$Z : \Sigma \times I \rightarrow \Sigma \times I$$


$$X_t : \Sigma \hookrightarrow \Sigma \times I$$



# Configuration space

$$Q_{EM} = C^\infty(\Sigma) \times \Omega^1(\Sigma) \times \text{Emb}_{\text{sp}}(\Sigma, M)$$


$$TQ_{EM} = TC^\infty(\Sigma) \times T\Omega^1(\Sigma) \times T\text{Emb}_{\text{sp}}(\Sigma, M)$$



# Configuration space

$$Q_{EM} = C^\infty(\Sigma) \times \Omega^1(\Sigma) \times \text{Emb}_{\text{sp}}(\Sigma, M)$$

$$TQ_{EM} = \mathbf{TC}^\infty(\Sigma) \times T\Omega^1(\Sigma) \times T\text{Emb}_{\text{sp}}(\Sigma, M)$$




# Configuration space

$$Q_{EM} = C^\infty(\Sigma) \times \Omega^1(\Sigma) \times \text{Emb}_{\text{sp}}(\Sigma, M)$$

$$TQ_{EM} = \mathbf{TC}^\infty(\Sigma) \times T\Omega^1(\Sigma) \times T\text{Emb}_{\text{sp}}(\Sigma, M)$$

$$\begin{matrix} \mathbf{\mathcal{C}}^\infty(\Sigma) \times \mathbf{\mathcal{C}}^\infty(\Sigma) \\ \Downarrow \end{matrix}$$




# Configuration space

$$Q_{EM} = C^\infty(\Sigma) \times \Omega^1(\Sigma) \times \text{Emb}_{\text{sp}}(\Sigma, M)$$

$$TQ_{EM} = TC^\infty(\Sigma) \times T\Omega^1(\Sigma) \times T\text{Emb}_{\text{sp}}(\Sigma, M)$$


$$\begin{array}{c} \mathcal{V} \\ || \\ C^\infty(\Sigma) \times C^\infty(\Sigma) \end{array}$$



# Configuration space

$$Q_{EM} = C^\infty(\Sigma) \times \Omega^1(\Sigma) \times \text{Emb}_{\text{sp}}(\Sigma, M)$$


$$TQ_{EM} = TC^\infty(\Sigma) \times T\Omega^1(\Sigma) \times T\text{Emb}_{\text{sp}}(\Sigma, M)$$
$$\begin{matrix} \parallel \\ C^\infty(\Sigma) \times C^\infty(\Sigma) \end{matrix} \qquad \begin{matrix} \parallel \\ \Omega^1(\Sigma) \times \Omega^1(\Sigma) \end{matrix}$$



# Configuration space

$$Q_{EM} = C^\infty(\Sigma) \times \Omega^1(\Sigma) \times \text{Emb}_{\text{sp}}(\Sigma, M)$$

$$TQ_{EM} = TC^\infty(\Sigma) \times T\Omega^1(\Sigma) \times \text{TEmb}_{\text{sp}}(\Sigma, M)$$
$$\begin{matrix} \mathcal{V} \\ \parallel \\ C^\infty(\Sigma) \times C^\infty(\Sigma) \end{matrix} \quad \quad \begin{matrix} \mathcal{V} \\ \parallel \\ \Omega^1(\Sigma) \times \Omega^1(\Sigma) \end{matrix}$$




# $T\text{Emb}(\Sigma, M)$

$$\begin{array}{ccc} & TM & \\ V_X \nearrow & \downarrow & \\ \Sigma & \xrightarrow[X]{} & M \end{array}$$


$$V_x \in T_x \text{Emb}(\Sigma, M)$$

$$\{X_\lambda\} \quad / \quad \begin{cases} X_0 = X \\ \dot{X}_0 = V_x \end{cases}$$



$$\hookrightarrow X_0$$

$$(\Sigma \times I, g)$$



# Back to the Hamiltonian

# Hamiltonian vector field

## Symplectic form

$$\left(FL_{EM}^P(\mathcal{D}), j^*\Omega\right) \xrightarrow{j} \left(T^*Q_{EM}, \Omega\right)$$

If  $Z \in \mathfrak{X}(FL_{EM}^P(\mathcal{D}))$  then  $Z = (q_\perp, q, X, p; Z_{q_\perp}, Z_q, \vec{Z}_X, Z_p)$

# Hamiltonian vector field

## Symplectic form

$$\left(FL_{EM}^P(\mathcal{D}), j^*\Omega\right) \xrightarrow{j} \left(T^*Q_{EM}, \Omega\right)$$

If  $Z \in \mathfrak{X}(FL_{EM}^P(\mathcal{D}))$  then  $Z = (q_\perp, q, X, p; Z_{q_\perp}, Z_q, \vec{Z}_X, Z_p)$

$$\iota_Z j^*\Omega = dH = 0$$

# Hamiltonian vector field

## Symplectic form

$$\left(FL_{EM}^P(\mathcal{D}), j^*\Omega\right) \xrightarrow{j} \left(T^*Q_{EM}, \Omega\right)$$

If  $Z \in \mathfrak{X}(FL_{EM}^P(\mathcal{D}))$  then  $Z = (q_\perp, q, X, p; Z_{q_\perp}, Z_q, \vec{Z}_X, Z_p)$

$$\iota_Z j^*\Omega = dH = 0$$

- $(Z_q)_a = \mathcal{L}_{z_X^T} q_a + \frac{p_a}{\sqrt{\gamma_X}} Z_X^\perp - d \left( q_\perp Z_X^\perp \right)$
- $Z_p^a = \mathcal{L}_{z_X^T} p^a + \sqrt{\gamma_X} \nabla_b \left( Z_X^\perp (dq)^{ba} \right)$
- $Z_X^\perp (\nabla_b p^b) = 0$
- $Z_X^T$  arbitrary
- $\left( Z_{q_\perp} - \mathcal{L}_{z_X^T} q_\perp + q_a \nabla^a Z_X^\perp \right) \nabla_b p^b = 0$

# Hamiltonian vector field

## Symplectic form

$$\left(FL_{EM}^P(\mathcal{D}), j^*\Omega\right) \xrightarrow{j} \left(T^*Q_{EM}, \Omega\right)$$

If  $Z \in \mathfrak{X}(FL_{EM}^P(\mathcal{D}))$  then  $Z = (q_\perp, q, X, p; Z_{q_\perp}, Z_q, \vec{Z}_X, Z_p)$

$$\iota_Z j^*\Omega = dH = 0$$

- $(Z_q)_a = \mathcal{L}_{Z_X^T} q_a + \frac{p_a}{\sqrt{\gamma_X}} Z_X^\perp - d(q_\perp Z_X^\perp)$
- $Z_p^a = \mathcal{L}_{Z_X^T} p^a + \sqrt{\gamma_X} \nabla_b (Z_X^\perp (dq)^{ba})$
- $Z_X^\perp (\nabla_b p^b) = 0$       •  $Z_X^T$  arbitrary
- $(Z_{q_\perp} - \mathcal{L}_{Z_X^T} q_\perp + q_a \nabla^a Z_X^\perp) \nabla_b p^b = 0$

# Hamiltonian vector field

## Symplectic form

$$\left(FL_{EM}^P(\mathcal{D}), j^*\Omega\right) \xrightarrow{j} \left(T^*Q_{EM}, \Omega\right)$$

If  $Z \in \mathfrak{X}(FL_{EM}^P(\mathcal{D}))$  then  $Z = (q_\perp, q, X, p; Z_{q_\perp}, Z_q, \vec{Z}_X, Z_p)$

$$\iota_Z j^*\Omega = dH = 0$$

- $(Z_q)_a = \mathcal{L}_{z_X^T} q_a$
- $Z_p^a = \mathcal{L}_{z_X^T} p^a$
- $Z_X^\perp = 0$
- $Z_X^T$  arbitrary
- $Z_{q_\perp} = \mathcal{L}_{z_X^T} q_\perp$

-  M. Bauer, P. Harms, P.W. Michor, *Almost local metrics on shape space of hypersurfaces in n-space*, SIAM J. Imaging Sci. (2012) 5.1.
-  J.F. Barbero, J. Margalef-Bentabol, E.J.S. Villaseñor, *Hamiltonian dynamics of the parametrized electromagnetic field*, Classical and Quantum Gravity (2016) 33 [arXiv: 1511.00826]
-  J.F. Barbero, J. Margalef-Bentabol, E.J.S. Villaseñor, *Hamiltonian dynamics of the parametrized electromagnetic field*, Classical and Quantum Gravity (2016) 33 [arXiv: 1511.00826]
-  K. Kuchař, *Geometry of hyperspace. I*, J. Math. Phys. (1976) 17.

# Thanks for your attention

-  M. Bauer, P. Harms, P.W. Michor, *Almost local metrics on shape space of hypersurfaces in n-space*, SIAM J. Imaging Sci. (2012) 5.1.
-  J.F. Barbero, J. Margalef-Bentabol, E.J.S. Villaseñor, *Hamiltonian dynamics of the parametrized electromagnetic field*, Classical and Quantum Gravity (2016) 33 [arXiv: 1511.00826]
-  J.F. Barbero, J. Margalef-Bentabol, E.J.S. Villaseñor, *Hamiltonian dynamics of the parametrized electromagnetic field*, Classical and Quantum Gravity (2016) 33 [arXiv: 1511.00826]
-  K. Kuchař, *Geometry of hyperspace. I*, J. Math. Phys. (1976) 17.