

Bigravitons as dark matter and gravitational waves

11th July 2016

Katsuki Aoki,

Waseda University (Japan)

Based on KA and Shinji Mukohyama,

“Massive gravitons as dark matter and gravitational waves”

PRD 94, 024001 (2016) [arXiv:1604.06704]

Introduction

GR can describe our Universe
if we introduce unknown matters.

We may need beyond standard model!!

Standard model of gravity = GR, but


What is graviton?

- It should be spin-2 field.
- Massless field or Massive field? How many gravitons?

There could be several gravitons as other gauge bosons.

For simplicity, we focus on the case of two gravitons

→ Bigravitons = massless graviton and massive graviton



Non-linear bigravity theory (Hassan, Rosen, '11)

Two dynamical metrics: $g_{\mu\nu}$ and $f_{\mu\nu}$ (Hassan, and Rosen, 2011)

$$S = \frac{1}{2\kappa_g^2} \int d^4x \sqrt{-g} R(g) + \frac{1}{2\kappa_f^2} \int d^4x \sqrt{-f} \mathcal{R}(f) \\ - \frac{m^2}{\kappa^2} \int d^4x \sqrt{-g} \sum_{i=0}^4 b_i \mathcal{U}_i(g, f) + S^{[m]}(g, \psi) \quad \kappa^2 = \kappa_g^2 + \kappa_f^2$$

$$\mathcal{U}_n(g, f) = -\frac{1}{n!(4-n)!} \epsilon^{\dots} \epsilon_{\dots} (\gamma^\mu{}_\nu)^n \quad \gamma^\mu{}_\alpha \gamma^\alpha{}_\nu = g^{\mu\alpha} f_{\alpha\nu}$$

Interaction is **restricted** to be free from ghost.

Particle aspect of massive graviton? = It may be just a massive particle.

We derive the energy-momentum (pseudo-)tensor of massive graviton from nonlinear bigravity theory.

→ **Massive graviton can be a dark matter.**

Linearization of bigravity

Consider the perturbation $\delta g_{\mu\nu} := g_{\mu\nu} - \eta_{\mu\nu}$, $\delta f_{\mu\nu} := f_{\mu\nu} - \eta_{\mu\nu}$.

Mass eigenstates with mass dimension one

$$h_{\mu\nu} := \frac{\kappa_f}{\kappa_g \kappa} \delta g_{\mu\nu} + \frac{\kappa_g}{\kappa_f \kappa} \delta f_{\mu\nu}, \quad \varphi_{\mu\nu} := \frac{1}{\kappa} (\delta g_{\mu\nu} - \delta f_{\mu\nu}),$$

Linearized bigravity theory

= General Relativity (massless spin-2) + Fierz-Pauli theory (massive spin-2)

$$\begin{aligned} S_2 &= \int d^4x \left[\frac{1}{\kappa_g^2} \mathcal{L}_{\text{EH}}[\delta g] + \frac{1}{\kappa_f^2} \mathcal{L}_{\text{EH}}[\delta f] + \mathcal{L}_{\text{int}}[\delta g, \delta f] + \frac{1}{2} \delta g_{\mu\nu} T_m^{\mu\nu} \right] \\ &= \int d^4x \left[\underbrace{\mathcal{L}_{\text{EH}}[h] + \frac{1}{2M_{\text{pl}}} h_{\mu\nu} T_m^{\mu\nu}}_{\text{GR (massless)} \quad M_{\text{pl}} := \frac{\kappa}{\kappa_g \kappa_f}} + \underbrace{\mathcal{L}_{\text{EH}}[\varphi] + \mathcal{L}_{\text{FP}}[\varphi] + \frac{1}{2M_G} \varphi_{\mu\nu} T_m^{\mu\nu}}_{\text{FP (massive)} \quad M_G := \frac{\kappa}{\kappa_g^2}} \right], \end{aligned}$$

Massive graviton as dark matter

We shall discuss gravitons perturbatively $\leftarrow T_{gw}^{\mu\nu}, T_G^{\mu\nu} = 2^{\text{nd}}$ order of EOM

We focus on sub-horizon scale

\rightarrow background can be Minkowski spacetime

Perturbative expansion can be used when

Vainshtein radius is much smaller than scales of DM

$\rightarrow p \ll \Lambda_3 \quad (\Lambda_3 := (mM_G)^{1/3})$

The universe is free from the Higuchi type instability

$\rightarrow H \lesssim m$


Energy-momentum tensor of graviton

In GR, $T_{gw}^{\mu\nu}$ can be obtained from the nonlinear part of EOM. (Isaacson, 1968)

$$T_{gw}^{\mu\nu} := -\langle M_{pl}^2 \delta G^{\mu\nu} \rangle = \frac{1}{4} \langle \partial^\mu h^{\alpha\beta} \partial^\nu h_{\alpha\beta} \rangle \text{ in transverse-traceless gauge}$$

→ (High-frequency) gravitons are sources of (low-frequency) graviton via nonlinear terms of Einstein equation in GR.

$$\mathcal{E}^{\mu\nu, \alpha\beta} h_{\alpha\beta} = \frac{1}{M_{pl}} (T_m^{\mu\nu} + T_{gw}^{\mu\nu})$$



Bigravity? There are two Einstein equations → Source of which equation?

Energy-momentum of gravitons in bigravity

We find the energy-momentum tensors (both massless and massive) are sources of the massless graviton.

The fully nonlinear Einstein equations in bigravity

$$G^{\mu\nu}(g) = \kappa_g^2 \left(T_g^{(\text{int})\mu\nu} + T_m^{\mu\nu} \right), \quad G^{\mu\nu}(f) = \kappa_f^2 T_f^{(\text{int})\mu\nu}$$

EOM of massless graviton including nonlinear terms:

$$\mathcal{E}^{\mu\nu,\alpha\beta} h_{\alpha\beta} = \frac{1}{M_{\text{pl}}} (T_m^{\mu\nu} + T_{\text{gw}}^{\mu\nu}(h) + T_G^{\mu\nu}(\varphi))$$

$$T_{\text{gw}}^{\mu\nu} := \left\langle -M_{\text{pl}}^2 \delta G^{(2)\mu\nu}(h) \right\rangle = \frac{1}{4} \langle \partial^\mu h^{\alpha\beta} \partial^\nu h_{\alpha\beta} \rangle$$

$$T_G^{\mu\nu} := \left\langle -\frac{1}{\kappa^2} \delta G^{(2)\mu\nu}(\varphi) + \delta T_g^{(2)(\text{int})\mu\nu}(h, \varphi) + \delta T_f^{(2)(\text{int})\mu\nu}(h, \varphi) \right\rangle = \frac{1}{4} \langle \partial^\mu \varphi^{\alpha\beta} \partial^\nu \varphi_{\alpha\beta} \rangle$$

Massive graviton as dark matter in bigravity

When we focus on the scales $\gg m^{-1}$,
the gravity mediated by the massive graviton is suppressed.

$$\Phi := -\delta g_{00}/2 = -\frac{GM}{r} (1 + \alpha e^{-mr}) \approx -\frac{GM}{r} \quad G := 1/8\pi M_{\text{pl}}^2$$

→ Only the Einstein equation of massless graviton is relevant.

$$\mathcal{E}^{\mu\nu,\alpha\beta} h_{\alpha\beta} = \frac{1}{M_{\text{pl}}} (T_{\text{m}}^{\mu\nu} + T_{\text{gw}}^{\mu\nu} + T_G^{\mu\nu}) \quad T_G^{\mu\nu} = \frac{1}{4} \langle \partial^\mu \varphi^{\alpha\beta} \partial^\nu \varphi_{\alpha\beta} \rangle$$

In the scales $\gg m^{-1}$,
the massive gravitons does not modify the gravity as a mediator,
however they can modify the gravity as a gravitational source.

When $m \gg p \gg \text{galaxy scales}^{-1}$ ($\Leftrightarrow m \gtrsim 10^{-23}$ eV)

$$T_G^{\mu\nu} = \frac{m^2}{4} \text{diag}[\langle \varphi^{\alpha\beta} \varphi_{\alpha\beta} \rangle, 0, 0, 0] = \text{Dark matter}$$

Other constraint on massive graviton DM

Lower bound on graviton mass: $m \gtrsim 10^{-23}$ eV

$$T_G^{\mu\nu} = \frac{m^2}{4} \text{diag}[\langle \varphi^{\alpha\beta} \varphi_{\alpha\beta} \rangle, 0, 0, 0] = \text{Dark matter}$$

We can obtain upper bound on graviton mass from the decay rate.


$$S_{\text{FP}} = \int d^4x \left[\mathcal{L}_{\text{EH}}[\varphi] + \mathcal{L}_{\text{FP}}[\varphi] + \frac{1}{2M_G} \varphi_{\mu\nu} T_m^{\mu\nu} \right]$$

Massive gravitons can decay to light particles.

Total decay rate of massive graviton

$$\Gamma_G \sim 0.1 \frac{m^3}{M_G^2} \ll H_0$$

Upper bound on graviton mass: $m \lesssim 0.01 \left(\frac{M_G}{M_{\text{pl}}} \right)^{2/3} \text{ GeV}$



Massive graviton production


The massive graviton can be dark matter when

$$10^{-23} \text{ eV} \lesssim m \lesssim 0.01 \left(\frac{M_G}{M_{\text{pl}}} \right)^{2/3} \text{ GeV},$$

We assume the production of massive graviton happens when $H_* \lesssim m$

How to generate massive gravitons?

$$\rightarrow \frac{1}{2M_{\text{pl}}} h_{\mu\nu} T_m^{\mu\nu} + \frac{1}{2M_G} \varphi_{\mu\nu} T_m^{\mu\nu}$$



When GWs are generated, MGs (=DM) are also generated.

→ **Bigravitons as dark matter and gravitational waves**

Massive gravitons from preheating

As an example, we consider generation through the preheating.

GW is generated by collisions of field bubbles (= large inhomogeneity) in sub-horizon scales.

GW momentum and energy $k_* \sim 1/R_*$, $\rho_{\text{gw}}^* \sim 0.1 (R_* H_*)^2 \rho_*$

R_* : typical size of field bubble

ρ_* : background energy density

Dufaux et al. JCAP 0903, 001 (2009)

Produced massive gravitons are relativistic ($\Lambda_3 \gg p^\mu \gg m$)

$$\langle \varphi_{\alpha\beta},{}^\mu \varphi^{\alpha\beta},{}^\nu \rangle \sim \frac{M_{\text{pl}}^2}{M_G^2} \langle h_{\alpha\beta},{}^\mu h^{\alpha\beta},{}^\nu \rangle$$

After the cosmic expansion, the present abundance is

$$\Omega_G \sim \frac{M_{\text{pl}}^2}{M_G^2} \frac{m}{2\pi f} \Omega_{\text{gw}} \quad f \text{ is the present frequency of GW}$$

Massive gravitons from preheating

We focus on GW to be sensitive in the LIGO range.

e.g. $\rho_*^{1/4} \sim 10^8 \text{ GeV}$, $R_*^{-1} \sim 0.1 \text{ GeV} \Rightarrow f \sim 40 \text{ Hz}$, $h^2 \Omega_{\text{gw}} \sim 10^{-9}$

If LIGO detects GW background and MG is dominant component of DM,

we obtain $\frac{M_{\text{pl}}^2}{M_G^2} m \sim 10^{-14} \text{ GeV}$

Consistency of our assumptions: $R_*^{-1} > m > \sqrt{2} H_*$

A set of consistent parameters is $m \sim 0.01 \text{ GeV}$, $M_G \sim 10^6 M_{\text{pl}}$

The corresponding free streaming scale $\sim 10^{-7} \text{ pc} \rightarrow \text{CDM}$

Even for our restricted case, we can construct a consistent model of DM.

Summary and Discussions

We have proposed a scenario in which **the massive graviton is DM**.

DM can be created when GM is created.

→ **GW can carry information about DM in our scenario.**

e.g. We can estimate the graviton mass from GW.

We construct a consistent model of DM originated by preheating.

→ $m \sim 0.01 \text{ GeV}$, $M_G \sim 10^6 M_{\text{pl}}$ when LIGO observes GW from preheating.

Direct detection of graviton mass? → consistency relation of MG-DM.

Other scenario would be possible.

Babichev et al. arXiv: 1604.08564.

✓ Thermal production → $\text{TeV} \lesssim m \lesssim 10^{11} \text{ GeV}$ with $M_G \gg M_{\text{pl}}$

✓ Other GW emission event? Inflation? First order transition?...