

Suppressing the primordial tensor amplitude without changing the scalar sector in quadratic curvature gravity

Kohji Yajima (Rikkyo University)

in collaboration with

Tsutomu Kobayashi (Rikkyo University)


Based on Phys. Rev. D **92**, 103503

[arXiv : 1508.07412]

GR21 @Columbia Univ.

July 13, 2016

Constraints on Inflation model



Planck 2015 results. XX
arXiv:1502.02114

Question

Can we modify only tensor modes
without changing the scalar sector?

Question

Can we modify only tensor modes
without changing the scalar sector?

Yes, we can
with higher curvature corrections

Construction of theories

Action: $S = S_{\text{EH}} + S_{\phi} + S_{\text{higher}}$

$$S_{\text{EH}} = \frac{1}{2\kappa} \int d^4x \sqrt{-g} \mathcal{R}, \quad \kappa = 8\pi G$$

$$S_{\phi} = \int d^4x \sqrt{-g} P(\phi, \partial^{\mu} \phi \partial_{\mu} \phi),$$

$$S_{\text{higher}} = \frac{1}{\kappa} \int d^4x \sqrt{-g} \left(\frac{1}{M^2} \mathcal{R}_{\mu\nu\rho\sigma} \mathcal{R}^{\mu\nu\rho\sigma} + \dots \right).$$

Construction of theories

Theories we want have the properties as follows:

- No ghost degrees of freedom
- Changing the dynamics of tensor perturbations while the scalar perturbations is left unchanged

Construction of theories

Construction with

the unit normal to constant ϕ hypersurfaces

$$u_\mu := -\frac{\partial_\mu \phi}{\sqrt{-\partial^\nu \phi \partial_\nu \phi}},$$

the induced metric

$$\gamma_{\mu\nu} = g_{\mu\nu} + u_\mu u_\nu,$$

for example: $\mathcal{R}_{\mu\nu\rho\sigma} \mathcal{R}_{\mu'\nu'\rho'\sigma'} \gamma^{\mu\mu'} \gamma^{\nu\nu'} \gamma^{\rho\rho'} u^\sigma u^{\sigma'}$

Construction of theories

ADM decomposition

taking constant ϕ hypersurfaces
as constant time hypersurfaces,

$$ds^2 = -N^2 dt^2 + \gamma_{ij} (dx^i + N^i dt) (dx^j + N^j dt) .$$

quadratic curvature terms

$$\sqrt{\gamma} N \times \{ K^4, K_{ij} K^{ij} K^2, \dots, R^2, R_{ij} R^{ij}, \\ K^2 R, K K^{ij} R_{ij}, \dots, D_i K_{jk} D^i K^{jk}, \dots \}$$

Cosmological perturbations

$$N = 1 + \delta N, \quad N_i = \partial_i \chi + \chi_i, \quad \gamma_{ij} = a^2 e^{2\zeta} (e^h)_{ij},$$

About scalar perturbations

$$K_i^j = H \delta_i^j + \frac{1}{3} \delta K \delta_i^j + \delta \tilde{K}_i^j,$$

where

$$\delta K = -3H\delta N + 3\dot{\zeta} - \frac{1}{a^2} \partial^2 \chi,$$

$$\delta \tilde{K}_i^j = -\frac{1}{a^2} \left(\partial_i \partial^j - \frac{1}{3} \delta_i^j \partial^2 \right) \chi,$$

and

$$\delta R_i^j = -\frac{1}{a^2} \left(\partial_i \partial^j + \delta_i^j \partial^2 \right) \zeta.$$

Combinations for which the scalar variables are canceled out

$$2\partial_i\delta\tilde{K}_{jk}\partial^i\delta\tilde{K}^{jk} - 3\partial_i\delta\tilde{K}^{ik}\partial^j\delta\tilde{K}_{jk},$$

and

$$\delta R_{ij}\delta R^{ij} - \frac{3}{8}\delta R^2,$$

Including vector and tensor perturbations

$$\begin{aligned} 2\partial_i\delta\tilde{K}_{jk}\partial^i\delta\tilde{K}^{jk} - 3\partial_i\delta\tilde{K}^{ik}\partial^j\delta\tilde{K}_{jk} \\ = \frac{1}{2a^2} \left(\partial_i \dot{h}_{jk} \right)^2 + \frac{1}{4a^6} \left(\partial^2 \chi_i \right)^2, \end{aligned}$$

$$\delta R_{ij}\delta R^{ij} - \frac{3}{8}\delta R^2 = \frac{1}{4a^4} \left(\partial^2 h_{ij} \right)^2.$$

Construction of Lagrangian

$$\mathcal{L}'_1 = \frac{\sqrt{\gamma}N}{M^2} \left(2D_i \tilde{K}_{jk} D^i \tilde{K}^{jk} - 3D_i \tilde{K}^{ik} D^j \tilde{K}_{jk} \right),$$

$$\mathcal{L}_2 = \frac{\sqrt{\gamma}N}{M^2} \left(R_{ij} R^{ij} - \frac{3}{8} R^2 \right),$$

As alternated for \mathcal{L}'_1

$$\mathcal{L}_1 = \frac{\sqrt{\gamma}N}{M^2} (2D_i \tilde{K}_{jk} D^i \tilde{K}^{jk} - D_i \tilde{K}^{ik} D^j \tilde{K}_{jk} - 2D_i \tilde{K}_{jk} D^j \tilde{K}^{ik})$$

this can be written as

$$\mathcal{L}_1 = \frac{\sqrt{\gamma}N}{M^2} W_{ijk} W^{ijk}, \quad W_{ijk} = 2D_{[i} \tilde{K}_{j]k} + D_l \tilde{K}^l_{[i} \gamma_{j]k}.$$

Construction of Lagrangian

$$\mathcal{L}'_1 = \frac{\sqrt{\gamma}N}{M^2} \left(2D_i \tilde{K}_{jk} D^i \tilde{K}^{jk} - 3D_i \tilde{K}^{ik} D^j \tilde{K}_{jk} \right),$$

$$\mathcal{L}_2 = \frac{\sqrt{\gamma}N}{M^2} \left(R_{ij} R^{ij} - \frac{3}{8} R^2 \right),$$

As alternated for \mathcal{L}'_1

$$\mathcal{L}_1 = \frac{\sqrt{\gamma}N}{M^2} (2D_i \tilde{K}_{jk} D^i \tilde{K}^{jk} - D_i \tilde{K}^{ik} D^j \tilde{K}_{jk} - 2D_i \tilde{K}_{jk} D^j \tilde{K}^{ik})$$

this can be written as

$$\mathcal{L}_1 = \frac{\sqrt{\gamma}N}{M^2} W_{ijk} W^{ijk}, \quad W_{ijk} = 2D_{[i} \tilde{K}_{j]k} + D_l \tilde{K}^l_{[i} \gamma_{j]k}.$$

$$= \frac{\sqrt{-g}}{M^2} C_{\mu\nu\rho\sigma} C_{\mu'\nu'\rho'\sigma'} \gamma^{\mu\mu'} \gamma^{\nu\nu'} \gamma^{\rho\rho'} u^\sigma u^{\sigma'}$$

Tensor amplitudes
in \mathcal{L}_1 and \mathcal{L}_2 model

\mathcal{L}_1 model

$$S = S_{\text{EH}} + S_{\phi} + S_{\text{higher}}$$

$$S_{\text{higher}} = \frac{1}{\kappa} \int d^4x \mathcal{L}_1$$

$$\mathcal{L}_1 = \frac{\sqrt{\gamma} N}{M^2} \left(2D_i \tilde{K}_{jk} D^i \tilde{K}^{jk} - D_i \tilde{K}^{ik} D^j \tilde{K}_{jk} - 2D_i \tilde{K}_{jk} D^j \tilde{K}^{ik} \right)$$

for tensor perturbations

$$S = \frac{1}{8\kappa} \int dt d^3x a^3 \left[\dot{h}_{ij}^2 - \frac{1}{a^2} (\partial_k h_{ij})^2 + \frac{4}{M^2 a^2} (\partial_k \dot{h}_{ij})^2 \right]$$

\mathcal{L}_1 model

$$f_k^\lambda(t) = \left(\frac{1}{4\kappa}\right)^{1/2} a^{3/2} \left(1 + \frac{4k^2}{M^2 a^2}\right)^{1/2} h_k^\lambda$$

$$\ddot{f}_k + \omega_k^2(t) f_k = 0$$

$$\omega_k^2 := -\frac{1}{4} \left(H^2 + 2\dot{H}\right) + \frac{k^2/a^2 - 2H^2 - \dot{H}}{1 + 4k^2/M^2 a^2} \\ - \frac{4H^2 k^2 / M^2 a^2}{(1 + 4k^2 / M^2 a^2)^2}$$

WKB solution for $k^2/a^2 \gg H^2, M^2$

$$f_k \simeq \frac{1}{\sqrt{2\omega_k}} \exp \left[-i \int^t \omega_k(t') dt' \right]$$

\mathcal{L}_1 model


$$\mathcal{P}_T(k) = \frac{k^3}{\pi^2} |h_k|^2$$

N. Deruelle et al. JHEP **09**, 009 (2012)

$$\mathcal{P}_T = \frac{2\kappa H^2}{\pi^2} \Xi_1(H/M),$$

where

$$\Xi_1(x) := \frac{\cosh(\pi\nu/2) \coth(\pi\nu/2) |\Gamma(-1/4 + i\nu/4)|^4}{128\pi^2 x^3},$$



$$\nu := \sqrt{x^{-2} - 1}$$

\mathcal{L}_2 model

$$S = S_{\text{EH}} + S_{\phi} + S_{\text{higher}}$$

$$S_{\text{higher}} = -\frac{1}{2\kappa} \int d^4x \mathcal{L}_2$$

$$\mathcal{L}_2 = \frac{\sqrt{\gamma}N}{M^2} \left(R_{ij}R^{ij} - \frac{3}{8}R^2 \right),$$

for tensor perturbations

$$S = \frac{1}{8\kappa} \int dt d^3x a^3 \left[\dot{h}_{ij}^2 - \frac{1}{a^2} (\partial_k h_{ij})^2 - \frac{1}{M^2 a^4} (\partial^2 h_{ij})^2 \right]$$

\mathcal{L}_2 model

$$v_k^\lambda := (4\kappa)^{-1/2} a h_k^\lambda$$

$$\frac{d^2 v_k}{d\eta^2} + \omega_k^2(\eta) v_k = 0$$

$$\omega_k^2 := k^2 + \frac{k^4}{M^2 a^2} - \frac{1}{a} \frac{d^2 a}{d\eta^2}$$

WKB solution

$$v_k \simeq \frac{1}{\sqrt{2\omega_k}} \exp \left[-i \int^\eta \omega_k(\eta') d\eta' \right]$$


solution at large k in de Sitter background

$$v_k = \frac{e^{-\pi/8x} W_{i/4x, 3/4}(-ixk^2\eta^2)}{(-2xk^2\eta)^{1/2}}$$

\mathcal{L}_2 model

$$\mathcal{P}_T = \frac{2\kappa H^2}{\pi^2} \Xi_2(H/M)$$

$$\Xi_2(x) := \frac{\pi}{4} \left[e^{\pi/(4x)} x^{3/2} |\Gamma(5/4 + i/(4x))|^2 \right]^{-1}$$



\mathcal{L}_2 model

\mathcal{L}_2 contains

$$\mathcal{L}_2 \sim \frac{1}{M^2} \zeta (\partial^2 \zeta)^2$$

non-Gaussianity generated by this term

$$f_{NL} \sim \frac{H^2}{\epsilon M^2}$$


$$f_{NL} \lesssim 1$$

$$\frac{H}{M} \lesssim \epsilon^{1/2} \ll 1$$


\mathcal{L}_2 model

$$\mathcal{P}_T = \frac{2\kappa H^2}{\pi^2} \Xi_2(H/M)$$


$$\Xi_2(x) := \frac{\pi}{4} \left[e^{\pi/(4x)} x^{3/2} |\Gamma(5/4 + i/(4x))|^2 \right]^{-1}$$



Suppression with \mathcal{L}_1 model



Suppression with \mathcal{L}_1 model




Summary

- We construct two possible theories which change only the dynamics of tensor perturbations without changing scalar sector.
- One of the theories, \mathcal{L}_1 , can decrease the tensor amplitude up to 65%.
- We can put some inflation models which are out of the observational constraints into the 2σ contour with this suppression effect.

\mathcal{L}_1 model

$$\mathcal{P}_T(k) = \frac{2\kappa H^2}{\pi^2} \Xi_1(H/M) \Big|_{k=aH}$$



Blue dashed line: analytic
Red points: numerical

Tensor to scalar ratio $r = 16\epsilon\Xi_1$


Tensor tilt $n_T := \mathrm{d} \ln \mathcal{P}_T / \mathrm{d} \ln k$

$$n_T = -\frac{2\epsilon}{1-\epsilon} \left[1 + \frac{1}{2} \frac{\mathrm{d} \ln \Xi_1}{\mathrm{d} \ln(H/M)} \right] \Big|_{k=aH}$$

Consistency relation

$$-8n_T/r \simeq \mathcal{D} \Big|_{k=aH}$$

$$\mathcal{D} := \frac{1 + (1/2)\mathrm{d} \ln \Xi_1 / \mathrm{d} \ln x}{\Xi_1} \Big|_{x=H/M}$$



Tensor to scalar ratio $r = 16\epsilon\Xi_1$

Tensor tilt $n_T := \mathrm{d} \ln \mathcal{P}_T / \mathrm{d} \ln k$

$$n_T = -\frac{2\epsilon}{1-\epsilon} \left[1 + \frac{1}{2} \frac{\mathrm{d} \ln \Xi_1}{\mathrm{d} \ln(H/M)} \right] \Big|_{k=aH} < 0$$

Consistency relation

$$-8n_T/r \simeq \mathcal{D} \Big|_{k=aH}$$

$$\mathcal{D} := \frac{1 + (1/2)\mathrm{d} \ln \Xi_1 / \mathrm{d} \ln x}{\Xi_1} \Big|_{x=H/M}$$

