

# Singularity Theorems in Regularity $C^{1,1}$

Michael Kunzinger

joint work with

R. Steinbauer

M. Stojković

J. A. Vickers

**21st International Conference  
on General Relativity and Gravitation**

# Singularity Theorems

# Singularity Theorems

Blueprint of the generic singularity theorem (J. Senovilla):

**Th.** If  $g$  is  $C^2$  then (i)–(iv) are incompatible, where

- (i) Energy condition.
- (ii) Causality condition.
- (iii) Initial or boundary condition.
- (iv) Causal completeness.

# Singularity Theorems

Blueprint of the generic singularity theorem (J. Senovilla):

**Th.** If  $g$  is  $C^2$  then (i)–(iv) are incompatible, where

- (i) Energy condition.
  - (ii) Causality condition.
  - (iii) Initial or boundary condition.
  - (iv) Causal completeness.
- $C^2$  is too much to ask! Realistic models (stars, matched spacetimes) involve jumps in matter variables  $\leadsto g \in C^{1,1}$ .

# Singularity Theorems

Blueprint of the generic singularity theorem (J. Senovilla):

**Th.** If  $g$  is  $C^2$  then (i)–(iv) are incompatible, where

- (i) Energy condition.
  - (ii) Causality condition.
  - (iii) Initial or boundary condition.
  - (iv) Causal completeness.
- $C^2$  is too much to ask! Realistic models (stars, matched spacetimes) involve jumps in matter variables  $\leadsto g \in C^{1,1}$ .
  - Theorem allows (i)–(iv) for  $C^{1,1}$ .

# Singularity Theorems

Blueprint of the generic singularity theorem (J. Senovilla):

**Th.** If  $g$  is  $C^2$  then (i)–(iv) are incompatible, where

- (i) Energy condition.
  - (ii) Causality condition.
  - (iii) Initial or boundary condition.
  - (iv) Causal completeness.
- $C^2$  is too much to ask! Realistic models (stars, matched spacetimes) involve jumps in matter variables  $\leadsto g \in C^{1,1}$ .
  - Theorem allows (i)–(iv) for  $C^{1,1}$ .
  - But  $C^{1,1}$  spacetimes are not singular (curvature bounded, unique geodesics).

# Singularity Theorems

Blueprint of the generic singularity theorem (J. Senovilla):

**Th.** If  $g$  is  $C^2$  then (i)–(iv) are incompatible, where

- (i) Energy condition.
  - (ii) Causality condition.
  - (iii) Initial or boundary condition.
  - (iv) Causal completeness.
- $C^2$  is too much to ask! Realistic models (stars, matched spacetimes) involve jumps in matter variables  $\leadsto g \in C^{1,1}$ .
  - Theorem allows (i)–(iv) for  $C^{1,1}$ .
  - But  $C^{1,1}$  spacetimes are not singular (curvature bounded, unique geodesics).
  - Below  $C^{1,1}$ : unbounded curvature, non-unique geodesics: singular.

# Singularity Theorems

Blueprint of the generic singularity theorem (J. Senovilla):

**Th.** If  $g$  is  $C^2$  then (i)–(iv) are incompatible, where

- (i) Energy condition.
  - (ii) Causality condition.
  - (iii) Initial or boundary condition.
  - (iv) Causal completeness.
- $C^2$  is too much to ask! Realistic models (stars, matched spacetimes) involve jumps in matter variables  $\leadsto g \in C^{1,1}$ .
  - Theorem allows (i)–(iv) for  $C^{1,1}$ .
  - But  $C^{1,1}$  spacetimes are not singular (curvature bounded, unique geodesics).
  - Below  $C^{1,1}$ : unbounded curvature, non-unique geodesics: singular.
  - Hence  $C^{1,1}$  is the natural threshold for singularity theorems.



# The Theorems of Penrose and Hawking

# The Theorems of Penrose and Hawking

## Penrose, 1965

Let  $(M, g)$  be a  $C^2$ - spacetime such that

- (i)  $\text{Ric}(X, X) \geq 0$  for every null vector  $X$ .
- (ii) There exists a non-compact Cauchy hypersurface  $S$  in  $M$
- (iii) There exists a compact achronal spacelike submanifold of codimension 2 with past-pointing timelike mean curvature vector field.

Then  $M$  is future null geodesically incomplete.

# The Theorems of Penrose and Hawking

## Penrose, 1965

Let  $(M, g)$  be a  $C^2$ -spacetime such that

- (i)  $\text{Ric}(X, X) \geq 0$  for every null vector  $X$ .
- (ii) There exists a non-compact Cauchy hypersurface  $S$  in  $M$
- (iii) There exists a compact achronal spacelike submanifold of codimension 2 with past-pointing timelike mean curvature vector field.

Then  $M$  is future null geodesically incomplete.

## Hawking, 1967

Let  $(M, g)$  be a  $C^2$ -spacetime such that

- (i)  $\text{Ric}(X, X) \geq 0$  for every timelike vector  $X$ .
- (ii) There exists a compact space-like hypersurface  $S$  in  $M$
- (iii) The unit normals to  $S$  are everywhere converging.

Then  $M$  is future time-like geodesically incomplete.

# The Theorems of Penrose and Hawking in $C^{1,1}$

## Penrose, 1965

Let  $(M, g)$  be a  $C^{1,1}$ -spacetime such that

- (i)  $\text{Ric}(X, X) \geq 0$  for every **Lip-cont. local null vector field**  $X$ .
- (ii) There exists a non-compact Cauchy hypersurface  $S$  in  $M$
- (iii) There exists a compact achronal spacelike submanifold of codimension 2 with past-pointing timelike mean curvature vector field.

Then  $M$  is future null geodesically incomplete.

## Hawking, 1967

Let  $(M, g)$  be a  $C^{1,1}$ -spacetime such that

- (i)  $\text{Ric}(X, X) \geq 0$  for every **smooth timelike local vector field**  $X$ .
- (ii) There exists a compact space-like hypersurface  $S$  in  $M$
- (iii) The unit normals to  $S$  are everywhere converging.

Then  $M$  is future time-like geodesically incomplete.

# Obstacles in the $\mathcal{C}^{1,1}$ -case – Strategy of proof

# Obstacles in the $\mathcal{C}^{1,1}$ -case – Strategy of proof

Main problems:

# Obstacles in the $\mathcal{C}^{1,1}$ -case – Strategy of proof

## Main problems:

- No appropriate version of Calculus of Variations available (second variation, maximizing curves, focal points, index form, ...)

# Obstacles in the $\mathcal{C}^{1,1}$ -case – Strategy of proof

## Main problems:

- No appropriate version of Calculus of Variations available (second variation, maximizing curves, focal points, index form, ...)
- $\mathcal{C}^2$ -causality theory rests on local equivalence with Minkowski space. This requires good properties of exponential map.
  - ▶  $\exp_p : \tilde{U} \rightarrow U$  homeomorphism.
  - ▶  $\exp_p(I^+(0) \cap \tilde{U}) = I^+(p) \cap U$ .
  - ▶ Radial geodesics are locally longest curves.



# Obstacles in the $\mathcal{C}^{1,1}$ -case – Strategy of proof

## Main problems:

- No appropriate version of Calculus of Variations available (second variation, maximizing curves, focal points, index form, ...)
- $\mathcal{C}^2$ -causality theory rests on local equivalence with Minkowski space. This requires good properties of exponential map.
  - ▶  $\exp_p : \tilde{U} \rightarrow U$  homeomorphism.
  - ▶  $\exp_p(I^+(0) \cap \tilde{U}) = I^+(p) \cap U$ .
  - ▶ Radial geodesics are locally longest curves.

## Strategy:

# Obstacles in the $\mathcal{C}^{1,1}$ -case – Strategy of proof

## Main problems:

- No appropriate version of Calculus of Variations available (second variation, maximizing curves, focal points, index form, ...)
- $\mathcal{C}^2$ -causality theory rests on local equivalence with Minkowski space. This requires good properties of exponential map.
  - ▶  $\exp_p : \tilde{U} \rightarrow U$  homeomorphism.
  - ▶  $\exp_p(I^+(0) \cap \tilde{U}) = I^+(p) \cap U$ .
  - ▶ Radial geodesics are locally longest curves.

## Strategy:

- $\exp_p$  is a bi-Lipschitz homeomorphism with good causal properties.

# Obstacles in the $\mathcal{C}^{1,1}$ -case – Strategy of proof

## Main problems:

- No appropriate version of Calculus of Variations available (second variation, maximizing curves, focal points, index form, ...)
- $\mathcal{C}^2$ -causality theory rests on local equivalence with Minkowski space. This requires good properties of exponential map.
  - ▶  $\exp_p : \tilde{U} \rightarrow U$  homeomorphism.
  - ▶  $\exp_p(I^+(0) \cap \tilde{U}) = I^+(p) \cap U$ .
  - ▶ Radial geodesics are locally longest curves.

## Strategy:

- $\exp_p$  is a bi-Lipschitz homeomorphism with good causal properties.
- Employ regularization adapted to causal structure following
  - ▶ Chrusciel/Grant:  $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$ ,  $\check{g}_\varepsilon, \hat{g}_\varepsilon \rightarrow g$  in  $C^1$ , curvatures loc. unif. bounded.

# Obstacles in the $\mathcal{C}^{1,1}$ -case – Strategy of proof

## Main problems:

- No appropriate version of Calculus of Variations available (second variation, maximizing curves, focal points, index form, ...)
- $\mathcal{C}^2$ -causality theory rests on local equivalence with Minkowski space. This requires good properties of exponential map.
  - ▶  $\exp_p : \tilde{U} \rightarrow U$  homeomorphism.
  - ▶  $\exp_p(I^+(0) \cap \tilde{U}) = I^+(p) \cap U$ .
  - ▶ Radial geodesics are locally longest curves.

## Strategy:

- $\exp_p$  is a bi-Lipschitz homeomorphism with good causal properties.
- Employ regularization adapted to causal structure following
  - ▶ Chrusciel/Grant:  $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$ ,  $\check{g}_\varepsilon, \hat{g}_\varepsilon \rightarrow g$  in  $C^1$ , curvatures loc. unif. bounded.
- Avoid Calculus of Variations.

# Obstacles in the $\mathcal{C}^{1,1}$ -case – Strategy of proof

## Main problems:

- No appropriate version of Calculus of Variations available (second variation, maximizing curves, focal points, index form, ...)
- $\mathcal{C}^2$ -causality theory rests on local equivalence with Minkowski space. This requires good properties of exponential map.
  - ▶  $\exp_p : \tilde{U} \rightarrow U$  homeomorphism.
  - ▶  $\exp_p(I^+(0) \cap \tilde{U}) = I^+(p) \cap U$ .
  - ▶ Radial geodesics are locally longest curves.


## Strategy:

- $\exp_p$  is a bi-Lipschitz homeomorphism with good causal properties.
- Employ regularization adapted to causal structure following  
▶ Chrusciel/Grant:  $\check{g}_\varepsilon \prec g \prec \hat{g}_\varepsilon$ ,  $\check{g}_\varepsilon, \hat{g}_\varepsilon \rightarrow g$  in  $C^1$ ,  
curvatures loc. unif. bounded.
- Avoid Calculus of Variations.
- Re-build causality theory for  $\mathcal{C}^{1,1}$ -metrics.

# The $\mathcal{C}^{1,1}$ -proof


# The $\mathcal{C}^{1,1}$ -proof

- $D^+(S) \subseteq D_{\check{g}_\varepsilon}^+(S)$ :



# The $\mathcal{C}^{1,1}$ -proof

- $D^+(S) \subseteq D_{\check{g}_\varepsilon}^+(S)$ :




- Limiting argument  $\Rightarrow$  for every  $p \in D^+(S)$  there exists a  $g$ -geodesic  $\gamma$  with  $L(\gamma) = d(S, p)$ .



# The $\mathcal{C}^{1,1}$ -proof


- $D^+(S) \subseteq D_{\tilde{g}_\varepsilon}^+(S)$ :



- Limiting argument  $\Rightarrow$  for every  $p \in D^+(S)$  there exists a  $g$ -geodesic  $\gamma$  with  $L(\gamma) = d(S, p)$ .
- $S$  compact  $\Rightarrow$  negative upper bound on  $\theta(0) = \lim \theta_\varepsilon(0)$ , where  $\theta$  trace of the extrinsic curvature of  $S$ .

# The $\mathcal{C}^{1,1}$ -proof


- $D^+(S) \subseteq D_{\tilde{g}_\varepsilon}^+(S)$ :



- Limiting argument  $\Rightarrow$  for every  $p \in D^+(S)$  there exists a  $g$ -geodesic  $\gamma$  with  $L(\gamma) = d(S, p)$ .
- $S$  compact  $\Rightarrow$  negative upper bound on  $\theta(0) = \lim \theta_\varepsilon(0)$ , where  $\theta$  trace of the extrinsic curvature of  $S$ .
- ▶ Regularize Ricci-curvature

# The $\mathcal{C}^{1,1}$ -proof


- $D^+(S) \subseteq D_{\check{g}_\varepsilon}^+(S)$ :



- Limiting argument  $\Rightarrow$  for every  $p \in D^+(S)$  there exists a  $g$ -geodesic  $\gamma$  with  $L(\gamma) = d(S, p)$ .
- $S$  compact  $\Rightarrow$  negative upper bound on  $\theta(0) = \lim \theta_\varepsilon(0)$ , where  $\theta$  trace of the extrinsic curvature of  $S$ .
- ▶ Regularize Ricci-curvature
- Ricci-curvature bound on  $\check{g}_\varepsilon$  and Raychaudhury equation  $\Rightarrow D^+(S)$  relatively compact, otherwise  $\exists$  focal points for  $\check{g}_\varepsilon$  too close to  $S$ .

## The $\mathcal{C}^{1,1}$ -proof


- $D^+(S) \subseteq D_{\check{g}_E}^+(S)$ :



- Limiting argument  $\Rightarrow$  for every  $p \in D^+(S)$  there exists a  $g$ -geodesic  $\gamma$  with  $L(\gamma) = d(S, p)$ .
- $S$  compact  $\Rightarrow$  negative upper bound on  $\theta(0) = \lim \theta_\varepsilon(0)$ , where  $\theta$  trace of the extrinsic curvature of  $S$ .
- ▶ Regularize Ricci-curvature
- Ricci-curvature bound on  $\check{g}_\varepsilon$  and Raychaudhuri equation  $\Rightarrow D^+(S)$  relatively compact, otherwise  $\exists$  focal points for  $\check{g}_\varepsilon$  too close to  $S$ .
- Therefore,  $H^+(S) \subseteq \overline{D^+(S)}$  compact.

## The $\mathcal{C}^{1,1}$ -proof

- $D^+(S) \subseteq D_{g_E}^+(S)$ :



- Limiting argument  $\Rightarrow$  for every  $p \in D^+(S)$  there exists a  $g$ -geodesic  $\gamma$  with  $L(\gamma) = d(S, p)$ .
- $S$  compact  $\Rightarrow$  negative upper bound on  $\theta(0) = \lim \theta_\varepsilon(0)$ , where  $\theta$  trace of the extrinsic curvature of  $S$ .
- ▶ Regularize Ricci-curvature
- Ricci-curvature bound on  $\check{g}_\varepsilon$  and Raychaudhuri equation  $\Rightarrow D^+(S)$  relatively compact, otherwise  $\exists$  focal points for  $\check{g}_\varepsilon$  too close to  $S$ .
- Therefore,  $H^+(S) \subseteq \overline{D^+(S)}$  compact.
- Derive a contradiction as in the  $\mathcal{C}^\infty$ -case.

# Outlook

# Outlook

## Related results, comparison geometry

- Riemannian comparison results: Ricci, mean curvature, Laplace, ...
- Grant (2011): Null-cone comparison theorems
- Grant/Treude (2013): Lorentzian volume comparison with model spaces (warped products), new proof of Hawking's theorem
- Graf (2016): Comparison geometry proof of  $C^{1,1}$ -Hawking theorem.
- Graf (2016): Rigidity results for singularity theorems.

# Outlook

## Related results, comparison geometry

- Riemannian comparison results: Ricci, mean curvature, Laplace, ...
- Grant (2011): Null-cone comparison theorems
- Grant/Treude (2013): Lorentzian volume comparison with model spaces (warped products), new proof of Hawking's theorem
- Graf (2016): Comparison geometry proof of  $C^{1,1}$ -Hawking theorem.
- Graf (2016): Rigidity results for singularity theorems.

## Current research


- Comparison approach to Penrose singularity theorem (Evolve trapped surface along null geodesics, quantify area), should also give new proof in  $C^{1,1}$ .
- Long term goal: ▶ Hawking-Penrose singularity theorem in  $C^{1,1}$ : Will require completely new methods.



# References


- P.T. Chrusciel, J.D.E. Grant, On Lorentzian causality with continuous metrics. *Classical Quantum Gravity* 29 (2012), no. 14, 145001, 32 pp.
- M. Graf, Volume comparison for  $C^{1,1}$ -metrics, *Ann. Glob. Anal. Geom.*, 2016.
- M. Graf, Splitting theorems for hypersurfaces, Preprint 2016.
- J.D.E. Grant, Areas and volumes for null cones. *Ann. Henri Poincaré* 12 (2011), no. 5, 965–985.
- S. W. Hawking, G.F.R. Ellis, *The Large Scale Structure of Space-Time*. Cambridge, 1973.
- S. W. Hawking, R. Penrose, *The singularities of gravitational collapse and cosmology*. *Proc. Roy. Soc. London Ser. A* 314 1970 529–548.
- M. Kunzinger, R. Steinbauer, M. Stojković. *The exponential map of a  $C^{1,1}$ -metric*. *Differential Geom. Appl.* 34 (2014), 14–24.
- M. Kunzinger, R. Steinbauer, M. Stojković, J.A. Vickers. *A regularisation approach to causality theory for  $C^{1,1}$ -Lorentzian metrics*. *Gen. Relativity Gravitation* 46 (2014), no. 8, 46:1738.
- M. Kunzinger, R. Steinbauer, M. Stojković, J.A. Vickers, *Hawking's singularity theorem for  $C^{1,1}$ -metrics*. *Classical and Quantum Gravity*, 32(7):075012, 2015.
- M. Kunzinger, R. Steinbauer, and J. A. Vickers, *The Penrose singularity theorem in regularity  $C^{1,1}$* . *Classical and Quantum Gravity* 32(15):155010, 2015.
- E. Minguzzi  
*Convex neighborhoods for Lipschitz connections and sprays*.  
*Monatsh. Math.* 177 (2015), no. 4, 569–625.
- R. Penrose, Gravitational collapse and space-time singularities.  
*Phys. Rev. Lett.* 14 1965 57–59.
- J. M. M. Senovilla, *Singularity theorems and their consequences*  
*Gen. Relativity Gravitation* 29, no. 5, 29:701 (1997)

# Chrusciel-Grant regularization of the metric




# Chrusciel-Grant regularization of the metric

- Locally,  $g_\varepsilon := g * \rho_\varepsilon$ , glued by partition of unity subordinate to  $U_i \subseteq M$ .




# Chrusciel-Grant regularization of the metric




- Locally,  $g_\epsilon := g * \rho_\epsilon$ , glued by partition of unity subordinate to  $U_i \subseteq M$ .
- Same way: construct time-like 1-form  $\omega$  s.t.  $|\omega(X)| \geq c_i > 0$  for all  $g$ -causal vf  $X$  with  $\|X\|_g = 1$ .

# Chrusciel-Grant regularization of the metric



- Locally,  $g_\epsilon := g * \rho_\epsilon$ , glued by partition of unity subordinate to  $U_i \subseteq M$ .
- Same way: construct time-like 1-form  $\omega$  s.t.  $|\omega(X)| \geq c_i > 0$  for all  $g$ -causal vf  $X$  with  $\|X\|_g = 1$ .
- $\check{g}_{\eta,\lambda} := g_\eta + \lambda \omega \otimes \omega$

# Chrusciel-Grant regularization of the metric



- Locally,  $g_\varepsilon := g * \rho_\varepsilon$ , glued by partition of unity subordinate to  $U_i \Subset M$ .
- Same way: construct time-like 1-form  $\omega$  s.t.  $|\omega(X)| \geq c_i > 0$  for all  $g$ -causal vf  $X$  with  $\|X\|_h = 1$ .
- $\check{g}_{\eta,\lambda} := g_\eta + \lambda\omega \otimes \omega$
- Adapt  $\lambda = \lambda(\varepsilon)$  and  $\eta = \eta(\varepsilon)$  locally s.t. for  $\varepsilon$  small

$$g(X, X) \leq 0 \wedge \|X\|_h = 1 \Rightarrow g_{\eta,\lambda}(X, X) < 0$$



# Regularization of Ricci-curvature



# Regularization of Ricci-curvature

**Th.**  $g$  a  $\mathcal{C}^{1,1}$ -metric,  $\text{Ric}(X, X) \geq 0$  for every  $\mathcal{C}^\infty$ -VF  $X$ ,  $K \subset\subset X$ . Then

$$\forall C > 0 \ \forall \delta > 0 \ \forall \kappa < 0 \ \forall \varepsilon \ \forall X \in TM|_K \text{ with } \check{g}_\varepsilon(X, X) \leq \kappa, \|X\|_h \leq C : \\ \text{Ric}_\varepsilon(X, X) > -\delta.$$

# Regularization of Ricci-curvature

**Th.**  $g$  a  $\mathcal{C}^{1,1}$ -metric,  $\text{Ric}(X, X) \geq 0$  for every  $\mathcal{C}^\infty$ -VF  $X$ ,  $K \subset\subset X$ . Then

$$\forall C > 0 \ \forall \delta > 0 \ \forall \kappa < 0 \ \forall \varepsilon \ \forall X \in TM|_K \text{ with } \check{g}_\varepsilon(X, X) \leq \kappa, \|X\|_h \leq C : \\ \text{Ric}_\varepsilon(X, X) > -\delta.$$

**Proof.**

- $\check{g}_\varepsilon - g * \rho_\varepsilon \rightarrow 0$  in  $\mathcal{C}^2 \rightsquigarrow$  suffices to consider  $g_\varepsilon := g * \rho_\varepsilon$ .
- $R_{jk} = R_{jki}^i = \partial_{x^i} \Gamma_{kj}^i - \partial_{x^k} \Gamma_{ij}^i + \Gamma_{im}^i \Gamma_{kj}^m - \Gamma_{km}^i \Gamma_{ij}^m$
- Blue terms $_{|\varepsilon}$  converge uniformly.
- For red terms use variant of Friedrich's Lemma:

$$(R_{jk} X^j X^k) * \rho_\varepsilon - R_{jk}^\varepsilon X^j X^k \rightarrow 0 \text{ uniformly}$$

$$\rho_\varepsilon \geq 0 \Rightarrow (R_{jk} X^j X^k) * \rho_\varepsilon \geq 0.$$



◀ Back

# The Hawking-Penrose Singularity Theorem

Hawking and Penrose, 1967

For a causally complete  $C^2$ -spacetime, the following can't all hold:

- 1 Every inextendible causal geodesic contains a pair of conjugate points,
- 2  $M$  contains no closed timelike curves and
- 3 there is a future or past trapped achronal set  $S$ .

## Corollary

$M$  must be causally incomplete if Einstein's equations hold and

- 1  $M$  contains no closed timelike curves.
- 2  $M$  satisfies an energy condition.
- 3 *Generality*: nontrivial curvature at some point of any causal geodesic.
- 4  $M$  contains either
  - ▶ a trapped surface
  - ▶ some  $p$  s.t. convergence of all null geodesics changes sign in the past
  - ▶ a compact spacelike hypersurface

[◀ Back](#)

# The Hawking-Penrose Singularity Theorem

Hawking and Penrose, 1967

For a causally complete  $C^2$ -spacetime, the following can't all hold:

- 1 Every inextendible causal geodesic contains a pair of conjugate points,
- 2  $M$  contains no closed timelike curves and
- 3 there is a future or past trapped achronal set  $S$ .

## Corollary

$M$  must be causally incomplete if Einstein's equations hold and

- 1  $M$  contains no closed timelike curves.
- 2  $M$  satisfies an energy condition.
- 3 *Generality*: nontrivial curvature at some point of any causal geodesic.
- 4  $M$  contains either
  - ▶ a trapped surface
  - ▶ some  $p$  s.t. convergence of all null geodesics changes sign in the past
  - ▶ a compact spacelike hypersurface

◀ Back