

Scattering for 2+1 Equivariant Einstein wave map system

Nishanth Gudapati

Yale

13 July 2016, Columbia

Wave Equation, Wave Maps

- Wave equation for a scalar $u : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$

$$\square_g u := -\partial_t^2 u + \partial_{x^1}^2 u + \partial_{x^2}^2 u = 0.$$

Wave Equation, Wave Maps

- Wave equation for a scalar $u : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$

$$\square_g u := -\partial_t^2 u + \partial_{x^1}^2 u + \partial_{x^2}^2 u = 0.$$

- Generalize to wave maps $U : (M, g) \rightarrow N, (N, h)$ a Riemannian manifold.

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0$$

Wave Equation, Wave Maps

- Wave equation for a scalar $u : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$

$$\square_g u := -\partial_t^2 u + \partial_{x^1}^2 u + \partial_{x^2}^2 u = 0.$$

- Generalize to wave maps $U : (M, g) \rightarrow N, (N, h)$ a Riemannian manifold.

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0$$

- A natural geometric generalization, $N = \mathbb{R}^n$ reduces to wave equation

Wave Equation, Wave Maps

- Wave equation for a scalar $u : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$

$$\square_g u := -\partial_t^2 u + \partial_{x^1}^2 u + \partial_{x^2}^2 u = 0.$$

- Generalize to wave maps $U : (M, g) \rightarrow N, (N, h)$ a Riemannian manifold.

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0$$

- A natural geometric generalization, $N = \mathbb{R}^n$ reduces to wave equation
- Semilinear hyperbolic 2nd order PDE (exhibits causality)

Wave Equation, Wave Maps

- Wave equation for a scalar $u : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$

$$\square_g u := -\partial_t^2 u + \partial_{x^1}^2 u + \partial_{x^2}^2 u = 0.$$

- Generalize to wave maps $U : (M, g) \rightarrow N, (N, h)$ a Riemannian manifold.

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0$$

- A natural geometric generalization, $N = \mathbb{R}^n$ reduces to wave equation
- Semilinear hyperbolic 2nd order PDE (exhibits causality)
- Satisfies the (strong) null condition

Wave Equation, Wave Maps

- Wave equation for a scalar $u : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$

$$\square_g u := -\partial_t^2 u + \partial_{x^1}^2 u + \partial_{x^2}^2 u = 0.$$

- Generalize to wave maps $U : (M, g) \rightarrow N, (N, h)$ a Riemannian manifold.

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0$$

- A natural geometric generalization, $N = \mathbb{R}^n$ reduces to wave equation
- Semilinear hyperbolic 2nd order PDE (exhibits causality)
- Satisfies the (strong) null condition
- H^1 data is critical.

The Cauchy Problem

Let Σ be the initial Cauchy surface and \mathbf{X} be its unit normal, then the Cauchy problem of wave maps is the following

$$\left. \begin{aligned} \square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\alpha\beta} \partial_\alpha U^j \partial_\beta U^k &= 0 \text{ on } M \\ U|_\Sigma &= U_0 \\ \mathbf{T}(U)|_\Sigma &= U_1 \end{aligned} \right\} \quad (1)$$

such that

$$\begin{aligned} U_0 &: \Sigma \rightarrow N \\ p &\rightarrow U_0(p) \end{aligned}$$

and

$$\begin{aligned} U_1 &: \Sigma \rightarrow T_{U_0} N \\ p &\rightarrow T_{U_0(p)} N \end{aligned}$$

for $p \in \Sigma$.

Previous Results in $M = \mathbb{R}^{m+1}$

Many deep, beautiful results.

Previous Results in $M = \mathbb{R}^{m+1}$

Many deep, beautiful results. Interplay of various fields of mathematics.
Non-exhaustive list here

Previous Results in $M = \mathbb{R}^{m+1}$

Many deep, beautiful results. Interplay of various fields of mathematics.
Non-exhaustive list here

- Christodoulou, Shatah, Tahvildar-Zadeh, Tao, Klainerman, Machedon, Selberg, Rodnianski, Raphael, Tataru, Sterbenz, Struwe, Nahmoud, Stefanov, Uhlenbeck, Merle, Kenig, Krieger, Schlag, Lawrie etc.

Critical Einstein-Wave Map System

Previous discussion was on a fixed background M , namely \mathbb{R}^{m+1} .

Critical Einstein-Wave Map System

Previous discussion was on a fixed background M , namely \mathbb{R}^{m+1} . Now consider a more general situation

$$\mathbf{E}_{\mu\nu} := \mathbf{R}_{\mu\nu} - \frac{1}{2}R_g g_{\mu\nu} = \alpha \mathbf{T}_{\mu\nu} \quad (2)$$

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0 \quad (3)$$

Critical Einstein-Wave Map System

Previous discussion was on a fixed background M , namely \mathbb{R}^{m+1} . Now consider a more general situation

$$\mathbf{E}_{\mu\nu} := \mathbf{R}_{\mu\nu} - \frac{1}{2}R_g g_{\mu\nu} = \alpha \mathbf{T}_{\mu\nu} \quad (2)$$

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0 \quad (3)$$

- \mathbf{R} Ricci tensor, \mathbf{E} Einstein tensor of (M, g)

Critical Einstein-Wave Map System

Previous discussion was on a fixed background M , namely \mathbb{R}^{m+1} . Now consider a more general situation

$$\mathbf{E}_{\mu\nu} := \mathbf{R}_{\mu\nu} - \frac{1}{2}R_g g_{\mu\nu} = \alpha \mathbf{T}_{\mu\nu} \quad (2)$$

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0 \quad (3)$$

- \mathbf{R} Ricci tensor, \mathbf{E} Einstein tensor of (M, g)
- Dynamical background (M, g)

Critical Einstein-Wave Map System

Previous discussion was on a fixed background M , namely \mathbb{R}^{m+1} . Now consider a more general situation

$$\mathbf{E}_{\mu\nu} := \mathbf{R}_{\mu\nu} - \frac{1}{2}R_g g_{\mu\nu} = \alpha \mathbf{T}_{\mu\nu} \quad (2)$$

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0 \quad (3)$$

- \mathbf{R} Ricci tensor, \mathbf{E} Einstein tensor of (M, g)
- Dynamical background (M, g)
- John Wheeler “Spacetime tells matter how to move; matter tells spacetime how to curve”

Cauchy Problem of Critical Einstein Equivariant Wave Map System

$$\left. \begin{aligned} \mathbf{E}_{\mu\nu} &= \alpha \mathbf{T}_{\mu\nu} && \text{on } M^{2+1} \\ \square_g u &= \frac{k^2 f_u(u) f(u)}{r^2} && \text{on } M^{2+1} \\ U|_{\Sigma} &= U_0 \\ T(U)|_{\Sigma} &= U_1 \end{aligned} \right\} \quad (4)$$

Cauchy Problem of Critical Einstein Equivariant Wave Map System

$$\left. \begin{aligned} \mathbf{E}_{\mu\nu} &= \alpha \mathbf{T}_{\mu\nu} && \text{on } M^{2+1} \\ \square_g u &= \frac{k^2 f_u(u) f(u)}{r^2} && \text{on } M^{2+1} \\ U|_{\Sigma} &= U_0 \\ T(U)|_{\Sigma} &= U_1 \end{aligned} \right\} \quad (4)$$

Initial data $(\Sigma, q, \mathbf{K}, U_0, U_1)$ smooth, compactly supported and satisfying constraint equations.

Cauchy Problem of Critical Einstein Equivariant Wave Map System

$$\left. \begin{aligned} \mathbf{E}_{\mu\nu} &= \alpha \mathbf{T}_{\mu\nu} && \text{on } M^{2+1} \\ \square_g u &= \frac{k^2 f_u(u) f(u)}{r^2} && \text{on } M^{2+1} \\ U|_{\Sigma} &= U_0 \\ T(U)|_{\Sigma} &= U_1 \end{aligned} \right\} \quad (4)$$

Initial data $(\Sigma, q, \mathbf{K}, U_0, U_1)$ smooth, compactly supported and satisfying constraint equations. In what follows assume k (homotopy degree) = 1

In double null-coordinates:

$$r^{-1}(2\partial_\xi Z \partial_\xi r - \partial_\xi^2 r) = \partial_\xi u \partial_\xi u \quad (5)$$

$$r^{-1} \partial_{\xi\eta}^2 r = \frac{e^{2Z}}{4} \frac{f^2(u)}{r^2} \quad (6)$$

$$r^{-1}(2\partial_\eta Z \partial_\eta r - \partial_\eta^2 r) = \partial_\eta u \partial_\eta u, \quad (7)$$

$$-4r^2 e^{-2Z} \partial_{\xi\eta}^2 Z = \frac{r^2}{2} \left(4e^{-2Z} \partial_\eta u \partial_\xi u + \frac{f^2(u)}{r^2} \right) \quad (8)$$

$$\square_{g(u)} u = \frac{f_u(u) f(u)}{r^2}. \quad (9)$$

Theorem

Suppose (M, g, U) is the globally hyperbolic, maximal development of the initial data of the 2+1 equivariant Einstein-wave map system (Σ_0, q_0, K_0) with $E_0 < \epsilon^2$ for ϵ sufficiently small, with

$$\int_0^u f(s) ds \rightarrow \infty \quad \text{as} \quad u \rightarrow \infty$$

then (M, g, U) is geodesically complete with global regularity.

Steps in the proof

Joint with Andersson and Szeftel

Steps in the proof

Joint with Andersson and Szeftel

- Set up and smallness from non-concentration of energy (PhD Thesis, 2014)

Steps in the proof

Joint with Andersson and Szeftel

- Set up and smallness from non-concentration of energy (PhD Thesis, 2014)
- Reduction to a 'semilinear' problem in null coordinates (blackboard)

Steps in the proof

Joint with Andersson and Szeftel

- Set up and smallness from non-concentration of energy (PhD Thesis, 2014)
- Reduction to a 'semilinear' problem in null coordinates (blackboard)
- Bootstrap assumption: $\sup_{Q_{\xi_b}} |\partial_{\xi} u| < C$, this implies $\sup_{Q_{\xi_b}} r^{-\delta} u < C, \delta \in (0, \frac{1}{2})$

Steps in the proof

Joint with Andersson and Szeftel

- Set up and smallness from non-concentration of energy (PhD Thesis, 2014)
- Reduction to a 'semilinear' problem in null coordinates (blackboard)
- Bootstrap assumption: $\sup_{Q_{\xi_b}} |\partial_{\xi} u| < C$, this implies $\sup_{Q_{\xi_b}} r^{-\delta} u < C, \delta \in (0, \frac{1}{2})$
- Improve this to $\delta = \frac{1}{2}$ using the fundamental solution

Steps in the proof

Joint with Andersson and Szeftel

- Set up and smallness from non-concentration of energy (PhD Thesis, 2014)
- Reduction to a 'semilinear' problem in null coordinates (blackboard)
- Bootstrap assumption: $\sup_{Q_{\xi_b}} |\partial_\xi u| < C$, this implies $\sup_{Q_{\xi_b}} r^{-\delta} u < C, \delta \in (0, \frac{1}{2})$
- Improve this to $\delta = \frac{1}{2}$ using the fundamental solution
- Apply the operator $D := \partial_R + \frac{1}{R}$ to the wave equation and use the fundamental solution again to obtain uniform bound for $\frac{u}{r}$

Steps in the proof

Joint with Andersson and Szeftel

- Set up and smallness from non-concentration of energy (PhD Thesis, 2014)
- Reduction to a 'semilinear' problem in null coordinates (blackboard)
- Bootstrap assumption: $\sup_{Q_{\xi_b}} |\partial_{\xi} u| < C$, this implies $\sup_{Q_{\xi_b}} r^{-\delta} u < C, \delta \in (0, \frac{1}{2})$
- Improve this to $\delta = \frac{1}{2}$ using the fundamental solution
- Apply the operator $D := \partial_R + \frac{1}{R}$ to the wave equation and use the fundamental solution again to obtain uniform bound for $\frac{u}{r}$
- Finally, transform the wave equation to a critical wave equation in $4 + 1$ dimensions for $v \sim \frac{u}{r}$

Steps in the proof

Joint with Andersson and Szeftel

- Set up and smallness from non-concentration of energy (PhD Thesis, 2014)
- Reduction to a 'semilinear' problem in null coordinates (blackboard)
- Bootstrap assumption: $\sup_{Q_{\xi_b}} |\partial_{\xi} u| < C$, this implies $\sup_{Q_{\xi_b}} r^{-\delta} u < C, \delta \in (0, \frac{1}{2})$
- Improve this to $\delta = \frac{1}{2}$ using the fundamental solution
- Apply the operator $D := \partial_R + \frac{1}{R}$ to the wave equation and use the fundamental solution again to obtain uniform bound for $\frac{u}{r}$
- Finally, transform the wave equation to a critical wave equation in $4 + 1$ dimensions for $v \sim \frac{u}{r}$
- Use energy estimates for the wave equation for v to prove global regularity.

Steps in the proof

Joint with Andersson and Szeftel

- Set up and smallness from non-concentration of energy (PhD Thesis, 2014)
- Reduction to a 'semilinear' problem in null coordinates (blackboard)
- Bootstrap assumption: $\sup_{Q_{\xi_b}} |\partial_\xi u| < C$, this implies $\sup_{Q_{\xi_b}} r^{-\delta} u < C, \delta \in (0, \frac{1}{2})$
- Improve this to $\delta = \frac{1}{2}$ using the fundamental solution
- Apply the operator $D := \partial_R + \frac{1}{R}$ to the wave equation and use the fundamental solution again to obtain uniform bound for $\frac{u}{r}$
- Finally, transform the wave equation to a critical wave equation in $4 + 1$ dimensions for $v \sim \frac{u}{r}$
- Use energy estimates for the wave equation for v to prove global regularity.
- Global regularity for u follows

Joint with Dodson.

- Morawetz estimate 1:

$$\int_{\mathbb{R}} \int_{\mathbb{R}^4} \frac{1}{|x|^3} v^2 \bar{\mu}_{\check{g}} \leq \|v_0\|_{\dot{H}^1(\mathbb{R}^4)} + \|v_1\|_{L^2(\mathbb{R}^4)} \quad (10)$$

- Morawetz estimate 2:

$$\int_{\mathbb{R}} \int_{|x| < \rho} \check{e} \bar{\mu}_{\check{g}} \leq \rho \left(\|v_0\|_{\dot{H}^1(\mathbb{R}^4)} + \|v_1\|_{L^2(\mathbb{R}^4)} \right). \quad (11)$$

Joint with Dodson.

- Morawetz estimate 1:

$$\int_{\mathbb{R}} \int_{\mathbb{R}^4} \frac{1}{|x|^3} v^2 \bar{\mu}_{\check{g}} \leq \|v_0\|_{\dot{H}^1(\mathbb{R}^4)} + \|v_1\|_{L^2(\mathbb{R}^4)} \quad (10)$$

- Morawetz estimate 2:

$$\int_{\mathbb{R}} \int_{|x| < \rho} \check{e} \bar{\mu}_{\check{g}} \leq \rho \left(\|v_0\|_{\dot{H}^1(\mathbb{R}^4)} + \|v_1\|_{L^2(\mathbb{R}^4)} \right). \quad (11)$$

-

$$\|v\|_X \leq \|v_0\|_{\dot{H}^1(\mathbb{R}^4)} + \|v_1\|_{L^2(\mathbb{R}^4)} + \|F\|_Y. \quad (12)$$

- Scattering:

$$\|v\|_{L^2 L^8(\mathbb{R} \times \mathbb{R}^4)} < \infty$$

Concluding remarks

- Stability of Minkowski for polarized case: Huneau (2014, 2015), based on decay estimates in harmonic coordinates (Lindblad-Rodnianski, 2004).

Concluding remarks

- Stability of Minkowski for polarized case: Huneau (2014, 2015), based on decay estimates in harmonic coordinates (Lindblad-Rodnianski, 2004).
- Blow-up profile, soliton solutions

Concluding remarks

- Stability of Minkowski for polarized case: Huneau (2014, 2015), based on decay estimates in harmonic coordinates (Lindblad-Rodnianski, 2004).
- Blow-up profile, soliton solutions
- Black hole stability problem

Concluding remarks

- Stability of Minkowski for polarized case: Huneau (2014, 2015), based on decay estimates in harmonic coordinates (Lindblad-Rodnianski, 2004).
- Blow-up profile, soliton solutions
- Black hole stability problem
- Super-critical EWM system.

Thank you, glad to be here