

Lighting Up Inspiring Binary Black Hole Systems



21st International Conference
on General Relativity
and Gravitation
Columbia University, New York

Session B1: Wed. July 13, 2016

Scott C. Noble (U. Tulsa)

M. Campanelli (RIT)

D. Bowen (RIT)

J. Krolik (JHU)

B. Mundim (Frankfurt U.)


H. Nakano (Kyoto U.)

M. Zilhao (Barcelona U.)

Y. Zlochower (RIT)

Based on:

- Noble++2012, Mundim++2014
- Zilhao & Noble 2014,
- Zilhao++2015, Noble++in-prep



Thanks to NSF PRAC OCI-0725070, NSF CDI AST-1028087, NSF ACI-1515969, NSF AST-1515982
Computing resources provided by NCSA/BlueWaters and TACC/Stampede

Multimessenger Synergy

Electromagnetic Surveys



Pan-STARRS:

- Running, 4 skies per month

Large Synoptic Survey Telescope (LSST):

- 2021-2032, one sky every 3 days

All-Sky Automated Survey for Supernovae (ASAS-SN)

- Running, 1 sky a night, not very sensitive;

Gravitational Wave Observatories



eLISA/NGO


~2034?



~2020

- GW Detection/Localization <---> EM Detection/Localization;
- GW and light are connected theoretically but originate in wholly different mechanisms
 - --> independently constrain models;
- Follow up (X-ray, sub-mm) observations can often be made via coordinated alert systems;
- Cosmological “Standard Sirens”: New Distance vs. Redshift Measurement
Schutz 1986, Chernoff+Finn 1993, Finn 1996, Holz & Hughes 2005

Strategy



Hopkins, Hernquist,
Di Matteo, Springel++

Farris++2014

Shi++2014

Noble++2012

Gold++2014

Matter:

Viscous Hydro.

MHD

MHD

MHD

Gravity:

Newtonian

Newtonian

Post-Newtonian

GR

Physical Time (not to scale)

Galactic Merger

Binary Formation

Inspiral

Merger

Re-equilibration

Newtonian Gravity

Numerical Relativity

Post-Newtonian

Static GR

Harm3d

Harm3d

Eulerian, high-resolution/shock-capturing, 3-d, ideal MHD, dynamical GR, HLL fluxes, parabolic reconstruction, dynamical FMR

DATA

DATA

Motivation of Simulation Design

Better Models

+ MHD

+ 3-d

+ GR


+ Radiative Cooling
+ Radiation Feedback

- MHD stresses \rightarrow Ang. Mom. transportation;
- Field dissipation and growth cannot be modeled w/ hydrodynamics;

- Buoyancy, and lasting turbulence only possible in 3-d.


- Post-Newtonian (PN) accuracy required for binary separations below $\sim 100M$;
- Significant mass can follow binary through much of this period (Noble++2012);
- NR needed for merger proper;
- Analytical metric provides freedom to grid for the gas instead of the spacetime. (Mundim++2014)

- Cooling provides a way to include more realistic thermodynamics consistent with its luminosity predictions;
 - No longer have to rely on $L \sim \dot{M}$;
- Eventually radiation feedback important in regions of non-smooth optical depths (e.g., “gap”)




- Mundim++2014

$t=56700.$



- “Excise” BBH to afford $O(100)$ orbits and arrive at relaxed disk;
- Will soon use for runs with resolved BH’s;
- Disk starts in “equilibrium”, threaded by poloidal magnetic field;


$t=56700.$



MHD Simulations Predict an EM Signature:

Noble++2012


Periodic Signal



$$\omega_{\text{peak}} = 2(\Omega_{\text{bin}} - \Omega_{\text{lump}})$$

Surface Density

$t=61200.$



(in frame co-rotating with lump)

Variability vs. Post-Newtonian Accuracy:

1.5PN


Same parameters
thinner initial disk

1.5PN-H


Different parameters
same initial thickness

2.5PN

Benchmark
(Noble++2012)



Top-down view of Surface Density




Less accurate metrics result in:

- Slightly weaker $m=1$ mode or over-density (lump) feature;
- Also, hints that thicker disks may weaken lump mode;

Variability vs. Post-Newtonian Accuracy:


1.5PN

Same parameters
thinner initial disk




1.5PN-H

Different parameters
same initial thickness



2.5PN

Benchmark
(Noble++2012)




- Differences between PN orders is no larger than small differences in initial data that help determine the disks's evolution;
- These differences provide us with a measure of the systematic error involved in our predictions;
- MHD turbulence \gg 2.5PN order terms at $a=20M$;

$q=1$

Mass Ratio Noble++in-prep

$q=1/2$








$q=1/5$

$q=1/10$

$q=1$

Mass Ratio Noble++in-prep

$q=1/2$



$q=1/5$


Top-down view of Surface Density

$q=1/10$

$q=1$

Mass Ratio Noble++in-prep

$q=1/2$



$q=1/5$

Top-down view of Surface Density

$q=1/10$

The Lump Puzzle

Noble++in-prep

- Lump is also seen in:
 - Newtonian MHD (Shi & Krolik 2012,2015);
 - 2-d viscous hydrodynamic simulations (e.g., D’Orazio++2012);
- Lump is coincident with degradation of “MRI Quality Factor” or resolution *within lump*;
 - Is it numerical? Do we lack the resolution to resolve the low B-field in the lump?
 - Is it artificial? Are we draining the region of sufficiently magnetized material?
 - Is it physical? Could the lump be a “dead zone” in which magnetic field is dissipated at a rate faster than can be brought in?

MRI Quality Factor:

$$Q^{(i)} = \frac{2\pi |b^i|}{\Delta x^{(i)} \Omega_K(r) \sqrt{\rho h + 2p_m}}.$$




Strategy:

Noble++2010, Hawley++2011,2013

- **“Bigger Disk”:**
 - Increase radial extent of the disk, keeping H/R the same;
 - Large extent increases reservoir of magnetic flux and mass;
- **“Injected Flux”:**
 - Magnetic flux from t=0 added to late-time snapshot of original run.
 - Net “vertical flux” can amplify other components and MRI.
 - Increases local magnetic energy density by only a few percent.
 - Emulates proposed explanations of state transitions in LXRB disks.


$t = 75600.$

Bigger Disk




Surface Density


Bigger Disk



Original



Flux-Injected






More magnetic flux led to:

- Without lump, less coherent temporal power spectrum, resembling more a slightly bent power law like those seen in single black hole disks.
- Richer spectral including more beat mode present with a larger reservoir.
- Periodic signals are present as long as there is a strong $m=1$ (lump) mode.


Bigger Disk

Original

Flux-Injected



Top-down view of Surface Density




More magnetic flux led to:

- Injected flux erased and prevented eventual development of $m=1$ mode.
- Bigger disk developed lump later, once surface density reached its steady state.

Bigger Disk

Original

Flux-Injected



$$\alpha_{\text{mag}} = \frac{\langle B^r B_\phi \rangle}{\langle p_{\text{mag}} \rangle} > 0.4 \Rightarrow \text{“Resolved MRI Turbulence”}$$


Guan & Gammie 2009, Sorathia++2012, Hawley++2011,2013

- Disks have resolved MRI throughout.
- No transition seen in “Bigger Disk” run across transition to the lump phase.
- Equivalent resolution in run without the lump.

Bigger Disk

Original

Flux-Injected



Average Specific Stress = $\frac{\langle B^r B_\phi \rangle}{\langle \rho \rangle} \lesssim 10^{-4} \Rightarrow$ Lump Formation

- From gradual redistribution of mass and flattening of surface density from initial conditions, stress per unit mass decreases.
- Transition to lump phase is evident, appears there is a threshold $\sim 1e-4$ below which lump forms;
- We are trying to understand original of the this threshold value.

Summary & Conclusions

- Our 3-d MHD, PN-regime simulations develop a high-Q signal that is non-trivially connected to the binary's orbit, but tied to the period of the beat mode between lump's orbit and the binary's orbit.
- The signal's strength degrades with decreasing mass ratio, implying that it can help diagnose properties of the binary, and it disappears altogether between $1/5 < q < 1/2$;
- At a separation of $20M$, with equal-mass binaries, differences in the metric at 1.5PN and 2.5PN orders are smaller than stochastic and systematic uncertainties, with PN-accuracy effects being even smaller for smaller mass ratios.
- Over density $m=1$ mode (lump), develops while disk's MRI is resolved and for different conditions, implying that the lump is physical and typical for similar disks.
- Beat signal is expected as long as the stress per unit mass is not too large, i.e. the disk is magnetically-submissive.

The Future is Bright!

**Stay-tuned to this channel for Dennis Bowen's talk on
minidisk dynamics next!**

Visualization by M. Vanmoer (NCSA)