


Accretion in Radiative Equipartition (AiRE) Disks

Yasaman K. Yazdi (work with N. Afshordi)
University of Waterloo and Perimeter Institute
July 14, 2016, GR21, NYC



Why We Need New Models

New models are necessary to better understand accretion disks.

- Thermal instability in current thin and slim disk models.
- Incomplete understanding of the source of viscosity.
- Incomplete understanding of disk spectra.
 - Spin estimates from Continuum Fitting and Fe Line methods are sometimes in tension.
 - Discrepancies between standard predictions for disk sizes and direct measurements.



Some well known analytic and semi-analytic disk solutions ¹




Figure: Locations approximately correspond to $\alpha = 0.1$ and $r = 20M$.

¹A. Sadowski, 2011, arXiv:1108.0396.

Disk physics described by four equations:

1) Conservation of Mass: $\dot{M} = -2\pi\Sigma\sqrt{\Delta}\frac{v}{\sqrt{1-v^2}}$

2) Conservation of angular momentum:

$$\frac{\dot{M}}{2\pi}(\mathcal{L} - \mathcal{L}_{in}) = -2hr^2 t_{r\phi}, \quad t_{r\phi} = \alpha p$$

3) Vertical equilibrium: $h^2\Omega^2 = \frac{p}{\rho}$

4) Energy balance: $Q^{vis} = Q^{rad}$

Thermally unstable because $Q^{vis} \propto T^8$ while $Q^{rad} \propto T^4$ in the inner regions.

AiRE disk solution to thermal instability: $P_{gas} = P_{rad}$ in inner region.

1) Conservation of Mass: $\dot{M} = -2\pi\Sigma\sqrt{\Delta}\frac{v}{\sqrt{1-v^2}}$

2) Conservation of angular momentum:

$$\frac{\dot{M}}{2\pi}(\mathcal{L} - \mathcal{L}_{in}) = -2hr^2 t_{r\phi}, \quad t_{r\phi} = \alpha p$$

3) Vertical equilibrium: $h^2\Omega^2 = \frac{p}{\rho}$

4) Radiative Equipartition: $P_{gas} = P_{rad}$




Figure: The equipartition radius, where the AiRE disk replaces the standard SS thin disk model.




Figure: Surface density of AiRE disks and SS disks for different black hole masses. $\dot{M}/\dot{M}_{Edd} = 0.1$ and $\alpha = 0.01$.

How Big Can a Black Hole Grow?²

Self-gravity becomes important in disks outside of some radius, r_{sg} . When $r_{sg} = r_{ISCO}$ there will no longer be any accretion.

Toomre criterion for a disk to be stable is $\frac{h\Omega^2}{\pi\Sigma} > 1$.




Figure: Upper mass limits for AiRE disks, with $\alpha = 0.1$ and $a = 0, 0.999$.

²A. King, arXiv:1511.08502.

Solve using the shooting or relaxation method.

$$\frac{dv}{dr} = \frac{\mathcal{N}(r, v)}{\mathcal{D}(r, v)} \quad (1)$$

Must continuously go from subsonic to supersonic flow at the sonic point. \mathcal{D} vanishes at the sonic point, so \mathcal{N} must also vanish there:

$$\mathcal{N}|_{r=r_{sonic}} = \mathcal{D}|_{r=r_{sonic}} = 0. \quad (2)$$

The value of \mathcal{L}_{in} must be chosen carefully to achieve this.

The topology of v differs for solutions with $\mathcal{L}_{in} > \mathcal{L}_{in}^t$ and $\mathcal{L}_{in} < \mathcal{L}_{in}^t$.

Sample Topology change of v with different values of \mathcal{L}_{in}





Figure: Topology change of v with different \mathcal{L}_{in} . The solution with correct inner boundary conditions lies at the transition point.

$\dot{M}/\dot{M}_{Edd} = 0.01$, and $\alpha = 0.1$.

Nature of the Sonic Point

The eigenvalues of the Jacobian $\mathcal{J} = \begin{pmatrix} \frac{\partial \mathcal{D}}{\partial r} & \frac{\partial \mathcal{D}}{\partial v} \\ \frac{\partial \mathcal{N}}{\partial r} & \frac{\partial \mathcal{N}}{\partial v} \end{pmatrix}$, tell us the nature of the sonic point. Same sign: nodal point. Opposite signs: saddle point. Imaginary: spiral (unphysical) point.

Conjecture: Saddle points are stable while Nodal points are unstable. Transition at $\alpha \sim 1.25 \left((1-a)^{2/5} (3+a) \right)^{1/3} \left(\frac{\dot{M}}{M^2} \right)^{1/24}$.



- AiRE disks are a solution to thermal instability in disks.
- AiRE disks modify disk properties such as radial velocity, surface temperature, and the Toomre parameter.
- Work is in progress to make precise the properties of AiRE disks such as their spectra.
- Preliminary results for maximum black hole masses and saddle to nodal transitions.