

Equatorial Orbits of Test Particles in a Conformastatic Background

Antonio C. Gutiérrez-Piñeres

Universidad Tecnológica de Bolívar, Cartagena (Colombia)

GR21. New York, July 11, 2016

Conformastationary spacetimes are
stationary spacetimes with a conformally flat space of orbits,

Conformastationary spacetimes are stationary spacetimes with a conformally flat space of orbits, and **conformastatic spacetimes** comprises the static subset .

(Synge, *Relativity: The General Theory*, 1960.)

Conformastationary spacetimes are stationary spacetimes with a conformally flat space of orbits, and **conformastatic spacetimes** comprises the static subset .

(Synge, *Relativity: The General Theory*, 1960.)

$$ds^2 = -c^2 e^{2\phi(r,z)} dt^2 + e^{-2\phi(r,z)} (dr^2 + dz^2 + r^2 d\varphi^2) .$$

Conformastatic line element:

$$ds^2 = -c^2 e^{2\phi(r,z)} dt^2 + e^{-2\phi(r,z)} (dr^2 + dz^2 + r^2 d\varphi^2)$$

Conformastatic line element:

$$ds^2 = -c^2 e^{2\phi(r,z)} dt^2 + e^{-2\phi(r,z)} (dr^2 + dz^2 + r^2 d\varphi^2)$$

+

Conformastatic line element:

$$ds^2 = -c^2 e^{2\phi(r,z)} dt^2 + e^{-2\phi(r,z)} (dr^2 + dz^2 + r^2 d\varphi^2)$$

+

Magnetic potential:

$$A_\alpha = (0, 0, 0, A_\varphi(r, z))$$

Conformastatic line element:

$$ds^2 = -c^2 e^{2\phi(r,z)} dt^2 + e^{-2\phi(r,z)} (dr^2 + dz^2 + r^2 d\varphi^2)$$

+

Magnetic potential:

$$A_\alpha = (0, 0, 0, A_\varphi(r, z))$$

=

Conformastatic line element:

$$ds^2 = -c^2 e^{2\phi(r,z)} dt^2 + e^{-2\phi(r,z)} (dr^2 + dz^2 + r^2 d\varphi^2)$$

+

Magnetic potential:

$$A_\alpha = (0, 0, 0, A_\varphi(r, z))$$

=

Magnetized conformastatic space-time

- *Conformastationary disk-haloes in Einstein-Maxwell gravity*
A.C. Gutiérrez-Piñeres,
Gen. Rel. Grav. 47 (2015) no.5, 54
- *Variational thermodynamics of relativistic thin disks*
A.C. Gutiérrez-Piñeres, C.S. Lopez-Monsalvo, H. Quevedo,
Gen. Rel. Grav. 47 (2015) no.12, 144
- *Exact Relativistic Magnetized Haloes around Rotating Disks* A.C. Gutiérrez-Piñeres, A.J.S. Capistrano,
Adv. Math. Phys. 2015 (2015) 916026
- *Conformastatic disk-haloes in Einstein-Maxwell gravity*
A.C. Gutiérrez-Piñeres, G.A. González, H. Quevedo,
Phys. Rev. D87 (2013) no.4, 044010

- *Conformastationary disk-haloes in Einstein-Maxwell gravity*
A.C. Gutiérrez-Piñeres,
Gen. Rel. Grav. 47 (2015) no.5, 54
- *Variational thermodynamics of relativistic thin disks*
A.C. Gutiérrez-Piñeres, C.S. Lopez-Monsalvo, H. Quevedo,
Gen. Rel. Grav. 47 (2015) no.12, 144
- *Exact Relativistic Magnetized Haloes around Rotating Disks* A.C. Gutiérrez-Piñeres, A.J.S. Capistrano,
Adv. Math. Phys. 2015 (2015) 916026
- *Conformastatic disk-haloes in Einstein-Maxwell gravity*
A.C. Gutiérrez-Piñeres, G.A. González, H. Quevedo,
Phys. Rev. D87 (2013) no.4, 044010
- *Test particles in a magnetized conformastatic spacetime*
A.C. Gutiérrez-Piñeres, A.J.S. Capistrano, H. Quevedo,
Phys. Rev. D93 (2016) no.12, 124009

Conserved quantities

$$\text{Energy } E : \quad p_t = -mce^{2\phi}\dot{t} \equiv -\frac{E}{c},$$

$$\text{Angular Momentum } L : \quad p_\varphi = mr^2e^{-2\phi}\dot{\varphi} + \frac{q}{c}A_\varphi \equiv L,$$

Conserved quantities

$$\text{Energy } E : \quad p_t = -mce^{2\phi}\dot{t} \equiv -\frac{E}{c},$$

$$\text{Angular Momentum } L : \quad p_\varphi = mr^2e^{-2\phi}\dot{\varphi} + \frac{q}{c}A_\varphi \equiv L,$$

$$p_\alpha p^\alpha = -\Sigma,$$

Conserved quantities

$$\text{Energy } E : \quad p_t = -mce^{2\phi}\dot{t} \equiv -\frac{E}{c},$$

$$\text{Angular Momentum } L : \quad p_\varphi = mr^2e^{-2\phi}\dot{\varphi} + \frac{q}{c}A_\varphi \equiv L,$$

$$p_\alpha p^\alpha = -\Sigma,$$

$$-c^2e^{2\phi}\dot{t}^2 + e^{-2\phi}(\dot{r}^2 + \dot{z}^2 + r^2\dot{\varphi}^2) = -\Sigma = \begin{cases} 0, & \text{null curves,} \\ -c^2, & \text{time-like curves,} \\ c^2, & \text{space-like curves.} \end{cases}$$

Conserved quantities

$$\text{Energy } E : p_t = -mce^{2\phi}\dot{t} \equiv -\frac{E}{c},$$

$$\text{Angular Momentum } L : p_\varphi = mr^2e^{-2\phi}\dot{\varphi} + \frac{q}{c}A_\varphi \equiv L,$$

$$p_\alpha p^\alpha = -\Sigma,$$

$$-c^2e^{2\phi}\dot{t}^2 + e^{-2\phi}(\dot{r}^2 + \dot{z}^2 + r^2\dot{\varphi}^2) = -\Sigma = \begin{cases} 0, & \text{null curves,} \\ -c^2, & \text{time-like curves,} \\ c^2, & \text{space-like curves.} \end{cases}$$

$$\text{The equatorial plane } (z = 0) : \dot{r}^2 + \Phi = \frac{E^2}{m^2c^2},$$

$$\text{Effective potential : } \Phi(r) \equiv \frac{L^2}{m^2r^2} \left(1 - \frac{qA_\varphi}{Lc}\right)^2 e^{4\phi} + \Sigma e^{2\phi}.$$

Circular Orbits

Conditions for the occurrence of circular orbits are

$$\frac{d\Phi}{dr} = 0, \quad \Phi = \frac{E^2}{m^2 c^2}. \quad (1)$$

Circular Orbits

Conditions for the occurrence of circular orbits are

$$\frac{d\Phi}{dr} = 0, \quad \Phi = \frac{E^2}{m^2 c^2}. \quad (1)$$

Angular momentum of the particle in circular motion:

$$L_{c\pm} = \frac{qA_\varphi}{c} + \frac{qrA_{\varphi,r}e^\phi \pm \sqrt{(qrA_{\varphi,r}e^\phi)^2 - 4\Sigma c^2 m^2 r^3 \phi_{,r} (2r\phi_{,r} - 1)}}{2ce^\phi (2r\phi_{,r} - 1)}. \quad (2)$$

Circular Orbits

Conditions for the occurrence of circular orbits are

$$\frac{d\Phi}{dr} = 0, \quad \Phi = \frac{E^2}{m^2 c^2}. \quad (1)$$

Angular momentum of the particle in circular motion:

$$L_{c\pm} = \frac{qA_\varphi}{c} + \frac{qrA_{\varphi,r}e^\phi \pm \sqrt{(qrA_{\varphi,r}e^\phi)^2 - 4\Sigma c^2 m^2 r^3 \phi_{,r} (2r\phi_{,r} - 1)}}{2ce^\phi (2r\phi_{,r} - 1)}. \quad (2)$$

Energy of the particle in circular motion:

$$E_{c\pm}^{(\pm)} = \pm mce^\phi \left(\Sigma + \xi_c^{(\pm)} \right)^{1/2}, \quad (3)$$

where

$$\xi_c^{(\pm)} = \frac{\left[qrA_{\varphi,r}e^\phi \pm \sqrt{(qrA_{\varphi,r}e^\phi)^2 - 4\Sigma c^2 m^2 r^3 \phi_{,r} (2r\phi_{,r} - 1)} \right]^2}{4m^2 c^2 r^2 (2r\phi_{,r} - 1)^2}.$$

Orbit in which the particle is located at rest
(r_r) as seen by an observer at infinity

$$L = 0, \quad \frac{d\Phi}{dr} = 0. \quad (4)$$

Orbit in which the particle is located at rest
 (r_r) as seen by an observer at infinity

$$L = 0, \quad \frac{d\Phi}{dr} = 0. \quad (4)$$

For an orbit with a rest radius $r = r_r$, the energy of the particle is,

$$E_{r\pm}^{(\pm)} = \pm mce^{\phi} \left(\Sigma + \xi_r^{(\pm)} \right)^{1/2} \quad (5)$$

where

$$\xi_r^{(\pm)} = \frac{q^2 e^{2\phi} [r A_{\varphi,r} \pm (r A_{\varphi,r} + 2A_{\varphi} (2r\phi_{,r} - 1))]^2}{4m^2 c^2 r^2 (2r\phi_{,r} - 1)^2}. \quad (6)$$

Last stable circular orbit

$$\frac{d\Phi}{dr} = 0, \quad \Phi = \frac{E^2}{m^2 c^2}, \quad \frac{d^2\Phi}{dr^2} = 0, \quad (7)$$

Last stable circular orbit

$$\frac{d\Phi}{dr} = 0, \quad \Phi = \frac{E^2}{m^2 c^2}, \quad \frac{d^2\Phi}{dr^2} = 0, \quad (7)$$

$$\begin{aligned} L_{lsc}^{\pm} = & \frac{qA_{\varphi}}{c} + \left\{ qe^{\phi} \left(8r^2 A_{\varphi,r} \phi_{,r}^2 + r^2 A_{\varphi,rr} - 4r A_{\varphi,r} \right) \right. \\ & \pm \left[q^2 e^{2\phi} \left(8r^2 A_{\varphi,r} \phi_{,r}^2 + r^2 A_{\varphi,rr} - 4r A_{\varphi,r} \right)^2 \right. \\ & - 4(q^2 r^2 A_{\varphi,r}^2 e^{2\phi} + 2c^2 \Sigma m^2 r^4 \phi_{,r}^2 + c^2 \Sigma m^2 r^4 \phi_{,rr}) \\ & \left. \left. \times (8r^2 \phi_{,r}^2 + 2r^2 \phi_{,rr} - 8r \phi_{,r} + 3) \right]^{1/2} \right\} \\ & \times \left[2ce^{\phi} (8r^2 \phi_{,r}^2 + 2r^2 \phi_{,rr} - 8r \phi_{,r} + 3) \right]^{-1} \end{aligned} \quad (8)$$

The field of a punctual mass in Einstein-Maxwell gravity

$$ds^2 = -c^2 e^{2\phi} dt^2 + e^{-2\phi} (dr^2 + dz^2 + r^2 d\varphi^2), \quad (9)$$

$$\begin{aligned} R_{\alpha\beta} - \frac{1}{2} g_{\alpha\beta} R &= k_0 E_{\alpha\beta}, \quad \nabla_\beta F^{\alpha\beta} = 0, \\ E_{\alpha\beta} &= \frac{1}{4\pi} \left\{ F_{\alpha\gamma} F_{\beta}{}^{\gamma} - \frac{1}{4} g_{\alpha\beta} F_{\gamma\delta} F^{\gamma\delta} \right\}, \\ F_{\alpha\beta} &= A_{\beta,\alpha} - A_{\alpha\beta}, \quad A_\alpha = (0, A_\varphi). \end{aligned} \quad (10)$$

A solution of the Einstein Maxwell equations

$$\phi(r, z) = \phi[U(r, z)], \quad (11)$$

$$\nabla^2 U(r, z) = 0. \quad (12)$$

A solution of the Einstein Maxwell equations

$$\phi(r, z) = \phi[U(r, z)], \quad (11)$$

$$\nabla^2 U(r, z) = 0. \quad (12)$$

$$\phi = -\ln(1 - U), \quad (13)$$

$$A_\varphi(r, z) = \frac{c^2}{G^{1/2}} \int_0^r \tilde{r} U(\tilde{r}, z) d\tilde{r}, \quad (14)$$

$$\nabla^2 U(r, z) = 0, \quad U(r, z) < 1.$$

A solution of the Einstein Maxwell equations

$$\phi(r, z) = \phi[U(r, z)], \quad (11)$$

$$\nabla^2 U(r, z) = 0. \quad (12)$$

$$\phi = -\ln(1 - U), \quad (13)$$

$$A_\varphi(r, z) = \frac{c^2}{G^{1/2}} \int_0^r \tilde{r} U(\tilde{r}, z) d\tilde{r}, \quad (14)$$

$$\nabla^2 U(r, z) = 0, \quad U(r, z) < 1.$$

$$B_r = \frac{c^2}{G^{1/2}} r U_{,r} \quad B_z = \frac{c^2}{G^{1/2}} r U_{,z}. \quad (15)$$

The field of a punctual mass in Einstein-Maxwell gravity

$$U(r, z) = -\frac{GM}{c^2 R}, \quad R^2 = r^2 + z^2, \quad (16)$$

The field of a punctual mass in Einstein-Maxwell gravity

$$U(r, z) = -\frac{GM}{c^2 R}, \quad R^2 = r^2 + z^2, \quad (16)$$

$$\phi(r, z) = -\ln \left(1 + \frac{GM}{c^2 R} \right), \quad (17)$$

The field of a punctual mass in Einstein-Maxwell gravity

$$U(r, z) = -\frac{GM}{c^2 R}, \quad R^2 = r^2 + z^2, \quad (16)$$

$$\phi(r, z) = -\ln \left(1 + \frac{GM}{c^2 R} \right), \quad (17)$$

$$A_\varphi(r, z) = \sqrt{GM} \left(1 - \frac{z}{R} \right), \quad (18)$$

The field of a punctual mass in Einstein-Maxwell gravity

$$U(r, z) = -\frac{GM}{c^2 R}, \quad R^2 = r^2 + z^2, \quad (16)$$

$$\phi(r, z) = -\ln \left(1 + \frac{GM}{c^2 R} \right), \quad (17)$$

$$A_\varphi(r, z) = \sqrt{G}M \left(1 - \frac{z}{R} \right), \quad (18)$$

$$B_r = G^{1/2} M r^2 R^{-3}, \quad B_z = G^{1/2} M r z R^{-3}. \quad (19)$$

The field of a punctual mass in Einstein-Maxwell gravity

$$U(r, z) = -\frac{GM}{c^2 R}, \quad R^2 = r^2 + z^2, \quad (16)$$

$$\phi(r, z) = -\ln \left(1 + \frac{GM}{c^2 R} \right), \quad (17)$$

$$A_\varphi(r, z) = \sqrt{GM} \left(1 - \frac{z}{R} \right), \quad (18)$$

$$B_r = G^{1/2} M r^2 R^{-3}, \quad B_z = G^{1/2} M r z R^{-3}. \quad (19)$$

$$\Phi(r) = \frac{c^6 r^2 (Lc - q\sqrt{GM})^2}{m^2 (c^2 r + GM)^4} + \frac{\Sigma c^4 r^2}{(c^2 r + GM)^2}. \quad (20)$$

Circular motion of a charged test particle

Circular motion of a charged test particle

The angular momentum for a circular orbit with radius r_c

$$L_{c\pm} = \frac{q\sqrt{GM}}{c} \mp \frac{(c^2 r_c + GM)m}{c^2} \sqrt{\frac{\Sigma GM}{c^2 r_c - GM}} \quad (21)$$

Circular motion of a charged test particle

The angular momentum for a circular orbit with radius r_c

$$L_{c\pm} = \frac{q\sqrt{GM}}{c} \mp \frac{(c^2 r_c + GM)m}{c^2} \sqrt{\frac{\Sigma GM}{c^2 r_c - GM}} \quad (21)$$

The energy for a circular orbit with radius r_c

$$E_{c\pm} = \pm \frac{mc^4}{(c^2 r_c + GM)} \sqrt{\frac{\Sigma r_c^3}{c^2 r_c - GM}} \quad (22)$$




Figure: Angular momentum of a neutral test particle in terms of the radius orbit r_c/M




Figure: Energy of a charged test particle in terms of the radius orbit r_c/M




Figure: Angular momentum of charged particles in a time-like circular orbit In this graphic the angular momentum L_{c+}^+/Mm is plotted as a function of the radius r_c/M for some values of q/m . The continuous curve corresponds to the value $q/m = 2\sqrt{2}$.

Charged test particle at rest ($L = 0$) respect to an observer at infinity


$$\frac{r_{r \pm}}{M} = \frac{c^2 q^2 - 2\Sigma G m^2 \pm \sqrt{c^2 q^2 (c^2 q^2 - 8\Sigma G m^2)}}{2\Sigma m^2 c^2}, \quad (23)$$

Charged test particle at rest ($L = 0$) respect to an observer at infinity


$$\frac{r_{r\pm}}{M} = \frac{c^2 q^2 - 2\Sigma G m^2 \pm \sqrt{c^2 q^2 (c^2 q^2 - 8\Sigma G m^2)}}{2\Sigma m^2 c^2}, \quad (23)$$

$$E_{r+}^{(\pm)} = + \frac{m c^2 \sqrt{2 q^2 \left[q^2 - 2 G m^2 \pm \sqrt{q^2 (q^2 - 8 G m^2)^2} \right]^3}}{\left[q^2 \pm \sqrt{q^2 (q^2 - 8 G m^2)^2} \right]^2},$$

$$E_{r-}^{(\pm)} = - E_{r+}^{(\pm)}. \quad (24)$$



(a)



(b)

Figure: Radii and Energy of the charged particles for the time-like orbits characterized by the conditions $L = 0$ and $d\Phi/dr = 0$.

The last stable circular orbit

The angular momentum of a particle in the last stable circular orbit

$$L_{lsc} \pm = \frac{q\sqrt{GM}}{c} \pm \frac{2\sqrt{2\Sigma}GMm}{c^2}. \quad (25)$$

The last stable circular orbit

The angular momentum of a particle in the last stable circular orbit

$$L_{lsc} \pm = \frac{q\sqrt{GM}}{c} \pm \frac{2\sqrt{2\Sigma}GMm}{c^2}. \quad (25)$$

The energy of a particle in the last stable circular orbit

$$E_{lsc} \pm = \pm \frac{3}{4} \sqrt{\frac{3\Sigma}{2}} mc. \quad (26)$$

The last stable circular orbit

For a null curve

$$L_{lsc\,o\,\pm} = q\sqrt{GM}/c. \quad (27)$$

The last stable circular orbit

For a null curve

$$L_{lsc0 \pm} = q\sqrt{GM}/c. \quad (27)$$

For a time-like particle it is

$$L_{lsc0 \pm} = (q\sqrt{G} \pm 2\sqrt{2Gm})M/c. \quad (28)$$

The last stable circular orbit

For a null curve

$$L_{lsc0 \pm} = q\sqrt{GM}/c. \quad (27)$$

For a time-like particle it is

$$L_{lsc0 \pm} = (q\sqrt{G} \pm 2\sqrt{2Gm})M/c. \quad (28)$$

If the charge of the particle is

$$q = 2\sqrt{2Gm}, \quad \text{then} \quad L_{lsc0-} = L_{c+} = L_r = 0. \quad (29)$$

The last stable circular orbit

For a null curve

$$L_{lsc0 \pm} = q\sqrt{GM}/c. \quad (27)$$

For a time-like particle it is

$$L_{lsc0 \pm} = (q\sqrt{G} \pm 2\sqrt{2Gm})M/c. \quad (28)$$

If the charge of the particle is

$$q = 2\sqrt{2Gm}, \quad \text{then} \quad L_{lsc0-} = L_{c+} = L_r = 0. \quad (29)$$

If the charge of the particle is

$$q = -2\sqrt{2Gm} \quad \text{then} \quad L_{lsc0+} = L_{c-} = L_r = 0. \quad (30)$$

The last stable circular orbit

For a null curve

$$L_{lsc0 \pm} = q\sqrt{GM}/c. \quad (27)$$

For a time-like particle it is

$$L_{lsc0 \pm} = (q\sqrt{G} \pm 2\sqrt{2Gm})M/c. \quad (28)$$

If the charge of the particle is

$$q = 2\sqrt{2Gm}, \quad \text{then} \quad L_{lsc0-} = L_{c+} = L_r = 0. \quad (29)$$

If the charge of the particle is

$$q = -2\sqrt{2Gm} \quad \text{then} \quad L_{lsc0+} = L_{c-} = L_r = 0. \quad (30)$$

The last stable circular orbit occurs at the radius $r = 3GM/c^2$, independently of the value of the charge. Moreover, on the last stable orbit the particle is at rest, if the value of the charge is $q = \pm 2\sqrt{2Gm}$.




Figure: Particle at the last stable circular orbit, black point, $(q/m = 2\sqrt{2}, r/M = 3, L_{lsc0+} = 4\sqrt{2}Mm)$. Particle at rest, white point, $(q/m = 2\sqrt{2}, r/M = 3, L_{lsc0-} = 0)$.

Perihelion advance in a conformastatic magnetized spacetime

Value of the perihelion advance for a neutral test mass in the gravitational field of a punctual mass, endowed with a magnetic field.

$$\Delta\varphi = \frac{5\pi GM}{c^2 r_c} . \quad (31)$$

Perihelion advance in a conformastatic magnetized spacetime

Value of the perihelion advance for a neutral test mass in the gravitational field of a punctual mass, endowed with a magnetic field.

$$\Delta\varphi = \frac{5\pi GM}{c^2 r_c} . \quad (31)$$

Value obtained for the Schwarzschild spacetime

$$\Delta\varphi = \frac{6\pi GM}{c^2 r_c} . \quad (32)$$

Perihelion advance in a conformastatic magnetized spacetime

Value of the perihelion advance for a neutral test mass in the gravitational field of a punctual mass, endowed with a magnetic field.

$$\Delta\varphi = \frac{5\pi GM}{c^2 r_c} . \quad (31)$$

Value obtained for the Schwarzschild spacetime

$$\Delta\varphi = \frac{6\pi GM}{c^2 r_c} . \quad (32)$$

The perihelion advance around a punctual magnetic mass is therefore always smaller than the value obtained in Einstein gravity alone. We conclude that the perihelion advance permits us to differentiate between a spherically symmetric mass and a conformally symmetric punctual magnetic mass.