

“**Gravitational**” scalar-tensor theory

Atsushi NARUKO (TiTech)

with : Daisuke Yoshida (TiTech) Shinji Mukohyama (YITP)

based on : CQG Letter (Apr. 2016) [1512.06977]

introduction

Einstein's General Relativity

- “**The classical theory of fields**” by Landau & Lifshitz,

GR is a **unique theory of gravity** provided

- composed only by metric and its derivatives
- 4D
- covariant theory
- EOM is (at most) 2nd order

- have to abandon one of assumptions above (or more ?)

metric ? > 4D ? covariance ? 2nd order EOM ?

f (R) theory

- The action is given by a **non-linear** function of **R**

- **EOM is 4th order** because $R \supset \ddot{g} + \dot{g}^2$

$$\int d^4x \sqrt{-g} f(R) \sim \int d^4x f(\ddot{g}) \rightarrow \dot{f}(\ddot{g}) \supset \ddot{\ddot{g}} : \text{EOM}$$

- Under a **Weyl (conformal) transformation**,

$$g_{\mu\nu} \rightarrow \Omega^2 g_{\mu\nu}$$

f (R) theory = Einstein + **canonical scalar**


➡ change of gravity law = introduction of matter !!

correspondence

metric + ϕ (scalar-tensor)
exotic matter ?

metric (purely gravitational)
modified gravity ?

canonical scalar



f(R) theory

Horndeski's theory

- the most general S-T theory **with 2nd order EOM for g & ϕ** :

canonical : $\mathcal{L} = X - V(\phi) = -\frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi - V(\phi)$

k-essence : $\mathcal{L}_2 = K(\phi, X)$ $X = -(\nabla\phi)^2/2$

KGB : $\mathcal{L}_3 = G(\phi, X)\square\phi$

Horndeski : $\mathcal{L}_4 = G_4(\phi, X) R - \frac{\partial G_4}{\partial X} [(\nabla\nabla\phi)^2 - (\square\phi)^2]$

$$\mathcal{L}_5 = G_5(\phi, X) G_{\mu\nu}\nabla^\mu\nabla^\nu\phi - \frac{1}{3}\frac{\partial G_5}{\partial X} [(\nabla\nabla\phi)^3 + \dots]$$

(cf. GLPV theory)

dual description ??

metric + ϕ (scalar-tensor)
exotic matter ?

metric (purely gravitational)
modified gravity ?

canonical scalar



f (R) theory

k-essence

?

KGB (\supset Box)

??

Horndeski (\supset Box²)

???

Horndeski (\supset Box³)

????



a special Horndeski



f (Gauss-Bonnet)



Kobayashi et.al. (2011)

question

Question

What if we introduce derivatives of R ? cf. $f(R)$

$$f\left(R, (\nabla R)^2, \square R, \dots\right)$$

What is the corresponding S-T theory, if it exists ??

model

model

- The action is given by **R** and **derivatives of R** :

$$f\left(R, (\nabla R)^2, \square R, \dots\right) \quad (\nabla R)^2 = g^{\mu\nu} \nabla_\mu R \nabla_\nu R$$

c.f. **f (R, \square R, ... , \square^n R)** theory Wands (1993)

f (Riemann) theory Deruelle et.al. (2009)

- We call this model “**gravitational** scalar-tensor theory” since the theory is constructed only in terms of **gravitational language**, namely metric & its derivatives.

Ostrogradski's ghost ??

- **non-degenerate** Lagrangian with **higher derivatives**

$$L = L(q, \dot{q}, \ddot{q}) \quad (d^2 L / d\dot{q}^2 \neq 0)$$

$$H = P_1 Q_2 + P_2 f(Q_1, Q_2, P_2) - L(Q_1, Q_2, P_2) \\ (Q_1 = q, Q_2 = \dot{q})$$

➡ **Hamiltonian is unbounded below**

- Although $f(R, (\nabla R)^2, \square R, \dots) \supset f(g, \dot{g}, \ddot{g}, \dots)$
there is **no Ostrogradsky's ghost** in our model
because the Lagrangian is **degenerate**.

proof of healthiness

- consider a simple model : $f\left(R, (\nabla R)^2\right)$
- rewrite **R** with **ϕ** & **λ** : **!! $\lambda \neq f_\phi$!!**

$$f\left(R, (\nabla R)^2\right) = f\left(\phi, (\nabla \phi)^2\right) - \lambda(\phi - R)$$
- conformal tr. : $g^{\mu\nu} = 2\lambda \tilde{g}^{\mu\nu}$

$$\frac{1}{2}\tilde{R} + f\left(\phi, 2\lambda (\tilde{\nabla} \phi)^2\right) - \frac{1}{2}(\tilde{\nabla} \chi)^2$$
 $\lambda \equiv e^{\sqrt{2/3} \chi}$
- 👉 # of d.o.f.s : **2 (GW) + 2 (scalar)** ($\neq 2 + 1$ in $f(R)$)
- 👉 sub class of bi-scalar k-essence \supset **healthy** domain

some remarks

- $(\Box R)^2$ is **not allowed** in general -> **Ostrogradski's ghost**
among models with the form of $f\left(R, (\nabla R)^2, \Box R\right)$

$$\Rightarrow K\left(R, (\nabla R)^2\right) + G\left(R, (\nabla R)^2\right) \times \Box R$$

- $(\Box R)^2$ **can be introduced** “with specific combinations”

$$Q\left(R, (\nabla R)^2\right) R + Q_X\left(R, (\nabla R)^2\right) \left[(\Box R)^2 - (\nabla_\mu \nabla_\nu R)^2\right]$$

- Way to construct a healthy gravitational S-T theory :

$$\mathcal{L}^{\text{Horn}}(\phi, g_{\mu\nu}) \rightarrow \mathcal{L}^{\text{Horn}}(R, g_{\mu\nu})$$

dual description ??

metric + ϕ (scalar-tensor)
exotic matter ?

metric (purely gravitational)
modified gravity ?

canonical scalar



f(R) theory

k-essence

?

KGB (\supset Box)

??

Horndeski (\supset Box²)

???

Horndeski (\supset Box³)

????

to be continued ...

summary

summary

- We have considered a theory of gravity in which the action is given by R and derivatives of R .
- Despite the higher derivative nature of action, the theory is healthy (if f is properly chosen)
= no ghost & no Ostrogradsky's instabilities
- # of d.o.f.s = 2 (GW) + 2 (scalar)
 \Leftrightarrow 2 scalars-tensor theory \supset bi-scalar k-essence
- Way to construct a healthy gravitational S-T theory :

$$\mathcal{L}^{\text{Horn}}(\phi, g_{\mu\nu}) \rightarrow \mathcal{L}^{\text{Horn}}(\textcolor{red}{R}, g_{\mu\nu})$$

Thank you very much
for your attention !!

ありがとう (Arigatou)

how to construct GST

Horndeski (# : **2** + **1**)

k-essence : $K(\phi, (\nabla\phi)^2)$

KGB : $G(\phi, X) \square\phi$

Horn. (L4) :

$$G_4 R - G_{4,X} \left[(\nabla\nabla\phi)^2 - (\square\phi)^2 \right]$$

Horn. (L5) :

$$G_5 G_{\mu\nu} \nabla^\mu \nabla^\nu \phi + \dots$$

Gravitational S-T (# : **2** + **2**)

$K(R, (\nabla R)^2)$

$G(R, X_R) \square R$

$$G_4 R - G_{4,X} \left[(\nabla\nabla R)^2 - (\square R)^2 \right]$$

$$G_5 \dots + \dots$$

how to construct GST

Horndeski (**2** + **1**)

gravitational S-T (**2** + **2**)

Horn : $\mathcal{L}_5 = G_5(\phi, X) G_{\mu\nu} \nabla^\mu \nabla^\nu \phi - \frac{1}{3} \frac{\partial G_5}{\partial X} \left[(\nabla \nabla \phi)^3 + \dots \right]$

$$\mathcal{L}_5 \supset f\left(R, (\nabla R)^2\right) \left[\nabla^\alpha (\nabla_\alpha \square - \square \nabla_\alpha) R - \frac{1}{2} (\nabla R)^2 - \frac{1}{2} R \square R \right]$$

$$+ \frac{1}{3} \frac{\partial f}{\partial (\nabla R)^2} \left(R, (\nabla R)^2 \right) \left[(\square R)^3 - 3(\square R) (\nabla_\mu \nabla_\nu R)^2 + 2(\nabla_\mu \nabla_\nu R)^3 \right]$$

identity : $G_{\mu\nu} \nabla^\mu \nabla^\nu \psi = \nabla (\nabla \square - \square \nabla) \psi - \frac{1}{2} \nabla R \nabla \psi - \frac{1}{2} R \square \psi$