

A group theoretic approach to shear-free radiating stars

G Z Abebe

Astrophysics and cosmology research unit
School of Maths, Stat and Comp Sc.
University of KwaZulu-Natal
Durban
South Africa

21st International Conference on General Relativity and Gravitation,
Columbia University,
New York, USA,
July 14, 2016

The spacetime

We consider the particular case of spherically symmetric, shear-free radiating stellar models. The line element for the interior for the spacetimes is given by

$$ds^2 = -A^2 dt^2 + B^2 [dr^2 + r^2 d\Omega^2], \quad (1)$$

where A and B are metric functions of t and r , and $d\Omega^2 \equiv d\theta^2 + \sin^2 \theta d\phi^2$. The acceleration and expansion are nonzero but the fluid is shear-free.

The energy momentum tensor has the form

$$T_{ab} = (\mu + p_{\perp}) u_a u_b + p_{\perp} g_{ab} + (p_{\parallel} - p_{\perp}) \chi_a \chi_b + q_a u_b + q_b u_a, \quad (2)$$

with heat flux and anisotropic stress. The fluid four-velocity $u^a = \frac{1}{A} \delta_0^a$ is comoving, χ^a is a unit four-vector along the radial direction ($u_a \chi^a = 0$), and the heat flow vector $q^a = (0, q, 0, 0)$ is radially directed ($u_a q^a = 0$).

The Einstein field equations for the heat conducting spherically symmetric anisotropic fluid (2) become

$$\mu = \frac{3}{A^2} \frac{B_t^2}{B^2} - \frac{1}{B^2} \left(2 \frac{B_{rr}}{B} - \frac{B_r^2}{B^2} + \frac{4B_r}{rB} \right), \quad (3a)$$

$$p_{\parallel} = \frac{1}{A^2} \left(-2 \frac{B_{tt}}{B} - \frac{B_t^2}{B^2} + 2 \frac{A_t}{A} \frac{B_t}{B} \right) + \frac{1}{B^2} \left(\frac{B_r^2}{B^2} + 2 \frac{A_r}{A} \frac{B_r}{B} + \frac{2}{r} \frac{A_r}{A} + \frac{2}{r} \frac{B_r}{B} \right), \quad (3b)$$

$$p_{\perp} = -\frac{2}{A^2} \frac{B_{tt}}{B} + 2 \frac{A_t}{A^3} \frac{B_t}{B} - \frac{1}{A^2} \frac{B_t^2}{B^2} + \frac{1}{r} \frac{A_r}{A} \frac{1}{B^2} + \frac{1}{r} \frac{B_r}{B^3} + \frac{A_{rr}}{A} \frac{1}{B^2} - \frac{B_r^2}{B^4} + \frac{B_{rr}}{B^3}, \quad (3c)$$

$$q = -\frac{2}{AB^2} \left(-\frac{B_{rt}}{B} + \frac{B_r B_t}{B^2} + \frac{A_r}{A} \frac{B_t}{B} \right), \quad (3d)$$

for the line element (1). The equations (3) describe the gravitational interactions in the interior of a shear-free spherically symmetric star with heat flux and anisotropic pressures.

The boundary of a radiating star divides the spacetime into interior and exterior regions. The interior spacetime (1) has to match across the boundary of the star to the Vaidya spacetime

$$ds^2 = - \left(1 - \frac{2m(v)}{R} \right) dv^2 - 2dv dR + R^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (4)$$

which is the exterior. Here the quantity $m(v)$ denotes the mass of the star as measured by an observer at infinity. Matching leads to the junction condition

$$(p_{\parallel})_{\Sigma} = (Bq)_{\Sigma}, \quad (5)$$

where the hypersurface Σ defines the boundary of the radiating sphere.

The boundary of a radiating star divides the spacetime into interior and exterior regions. The interior spacetime (1) has to match across the boundary of the star to the Vaidya spacetime

$$ds^2 = - \left(1 - \frac{2m(v)}{R} \right) dv^2 - 2dv dR + R^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (4)$$

which is the exterior. Here the quantity $m(v)$ denotes the mass of the star as measured by an observer at infinity. Matching leads to the junction condition

$$(p_{\parallel})_{\Sigma} = (Bq)_{\Sigma}, \quad (5)$$

where the hypersurface Σ defines the boundary of the radiating sphere. The junction condition (5) together with the field equations (2) gives

$$\begin{aligned} & 2\frac{B_{rt}}{AB^2} + 2\frac{B_{tt}}{A^2B} - 2\frac{A_t B_t}{A^3B} - 2\frac{B_r B_t}{AB^3} - 2\frac{A_r B_r}{AB^3} - 2\frac{A_r B_t}{A^2B^2} - \frac{B_r^2}{B^4} \\ & + \frac{B_t^2}{A^2B^2} - 2\frac{A_r}{rAB^2} - 2\frac{B_r}{rB^3} = 0, \end{aligned} \quad (6)$$

valid at the boundary of a shear-free radiating star.

Lie symmetry analysis

Gen Relativ Gravit (2012) 44:147–156
DOI 10.1007/s10714-011-1268-5

RESEARCH ARTICLE

A general class of Euclidean stars

K. S. Govinder · M. Govender

Received: 6 May 2011 / Accepted: 25 August 2011 / Published online: 9 September 2011
© Springer Science+Business Media, LLC 2011

Gen Relativ Gravit (2012) 44:147–156
DOI 10.1007/s10714-011-1268-5

RESEARCH ARTICLE

A general class of Euclidean stars

K. S. Govinder · M. Govender

Received: 6 May 2011 / Accepted: 25 August 2011 / Published online: 9 September 2011
© Springer Science+Business Media, LLC 2011

Int J Theor Phys (2013) 52:3244–3254
DOI 10.1007/s10773-013-1620-5

Lie Symmetries for a Conformally Flat Radiating Star

G.Z. Abebe · K.S. Govinder · S.D. Maharaj

Received: 16 October 2012 / Accepted: 13 April 2013 / Published online: 20 April 2013

Geodesic models generated by Lie symmetries

G. Z. Abebe · S. D. Maharaj · K. S. Govinder

Received: 2 July 2013 / Accepted: 2 December 2013 / Published online: 18 December 2013
© Springer Science+Business Media New York 2013

Geodesic models generated by Lie symmetries

G. Z. Abebe · S. D. Maharaj · K. S. Govinder

Received: 2 July 2013 / Accepted: 2 December 2013 / Published online: 18 December 2013
© Springer Science+Business Media New York 2013

Gen Relativ Gravit (2014) 46:1733
DOI 10.1007/s10714-014-1733-z

RESEARCH ARTICLE

Generalized Euclidean stars with equation of state

G. Z. Abebe · S. D. Maharaj · K. S. Govinder

Received: 4 December 2013 / Accepted: 11 April 2014 / Published online: 24 April 2014
© Springer Science+Business Media New York 2014



A group theoretic approach to shear-free radiating stars

G. Z. Abebe^a, S. D. Maharaj^b, K. S. Govinder^c

Astrophysics and Cosmology Research Unit, School of Mathematics, Statistics and Computer Science,
University of KwaZulu-Natal, Private Bag 54001, Durban 4000, South Africa

Received: 5 August 2015 / Accepted: 7 October 2015 / Published online: 19 October 2015

© The Author(s) 2015. This article is published with open access at Springerlink.com

A group theoretic approach to shear-free radiating stars

G. Z. Abebe^a, S. D. Maharaj^b, K. S. Govinder^c

Astrophysics and Cosmology Research Unit, School of Mathematics, Statistics and Computer Science,
University of KwaZulu-Natal, Private Bag 54001, Durban 4000, South Africa

Received: 5 August 2015 / Accepted: 7 October 2015 / Published online: 19 October 2015
© The Author(s) 2015. This article is published with open access at Springerlink.com

Radiating stars in the presence of electric charge

G.Z. Abebe^{1,*} and S.D. Maharaj^{1,†}

¹*Astrophysics and Cosmology Research Unit, School of Mathematics,
Statistics and Computer Science, University of KwaZulu-Natal,
Private Bag X54001, Durban 4000, South Africa*

We consider the general model of an accelerating, expanding and shearing radiating star in the presence of charge. Using a new set of variables arising from the Lie symmetries of differential equations we transform the boundary equation into ordinary differential equations. We present several new exact solutions for the master equation. A particular family of solution may be interpreted as a charged generalised Euclidean stars. This family admits a linear barotropic equation of state. In the uncharged limit we regain stellar models where proper and areal radii are equal, and its generalisation.

An n th order differential equation

$$F(r, t, A, B, A_r, B_r, A_t, B_t, A_{rr}, B_{rr}, A_{rt}, B_{rt}, A_{tt}, B_{tt}, \dots) = 0 \quad (7)$$

where $A = A(r, t)$ and $B = B(r, t)$, admits a Lie symmetry of the form

$$\begin{aligned} G = & \xi_1(r, t, A, B) \frac{\partial}{\partial r} + \xi_2(r, t, A, B) \frac{\partial}{\partial t} \\ & + \eta_1(r, t, A, B) \frac{\partial}{\partial A} + \eta_2(r, t, A, B) \frac{\partial}{\partial B} \end{aligned} \quad (8)$$

provided that

$$G^{[n]}F \Big|_{F=0} = 0 \quad (9)$$

where $G^{[n]}$ is the n th prolongation of the symmetry G . The process is algorithmic and so can be implemented by computer algebraic packages.

Using PROGRAM LIE (Head 1993), we find the set of symmetries

$$G_1 = -Af'(t)\frac{\partial}{\partial A} + f(t)\frac{\partial}{\partial t}, \quad (10a)$$

$$G_2 = A\frac{\partial}{\partial A} + B\frac{\partial}{\partial B}, \quad (10b)$$

$$G_3 = A\frac{\partial}{\partial A} + r\frac{\partial}{\partial r}, \quad (10c)$$

where $f(t)$ is an arbitrary function of t .

Using PROGRAM LIE (Head 1993), we find the set of symmetries

$$G_1 = -Af'(t)\frac{\partial}{\partial A} + f(t)\frac{\partial}{\partial t}, \quad (10a)$$

$$G_2 = A\frac{\partial}{\partial A} + B\frac{\partial}{\partial B}, \quad (10b)$$

$$G_3 = A\frac{\partial}{\partial A} + r\frac{\partial}{\partial r}, \quad (10c)$$

where $f(t)$ is an arbitrary function of t . We take a general linear combination

$$aG_1 + bG_2 + cG_3 = [c + b - af'(t)]A\frac{\partial}{\partial A} + bB\frac{\partial}{\partial B} + af(t)\frac{\partial}{\partial t} + cr\frac{\partial}{\partial r}, \quad (11)$$

to reduce the partial differential equation (6) into ordinary differential equations for further analysis. Note that a , b and c are arbitrary constants.

Then (11) gives the invariants

$$x = \frac{\exp\left(\int^t \frac{dt}{af(t)}\right)}{r^{1/c}}, \quad (12a)$$

$$A = \frac{h(x)}{f(t)} \exp\left(\int^t \frac{cdt}{af(t)} + \int^t \frac{bdt}{af(t)}\right), \quad (12b)$$

$$B = g(x)r^{b/c}, \quad (12c)$$

where $a \neq 0$ and $c \neq 0$. Note that g and h being arbitrary functions of x .

Then (11) gives the invariants

$$x = \frac{\exp\left(\int^t \frac{dt}{af(t)}\right)}{r^{1/c}}, \quad (12a)$$

$$A = \frac{h(x)}{f(t)} \exp\left(\int^t \frac{cdt}{af(t)} + \int^t \frac{bdt}{af(t)}\right), \quad (12b)$$

$$B = g(x)r^{b/c}, \quad (12c)$$

where $a \neq 0$ and $c \neq 0$. Note that g and h being arbitrary functions of x . Using the invariants (12) we can write (6) in the form

$$\begin{aligned} & \left[2a^2gx^{2b+2c+1}((b+c)g - xg')\right] h'h^2 + 2acx^{b+c+2}g^2g'h'h \\ & - 2c^2x^2g^3g'h' + [c^2xg^2(xg'^2 - 2g((b+c-1)g' - xg''))] h \\ & - \left[2acgx^{b+c+1}(g(xg'' + g') - xg'^2)\right] h^2 + [a^2x^{2(b+c)}(xg' - bg) \\ & \times ((b+2c)g - xg')] h^3 = 0, \end{aligned} \quad (13)$$

where primes denote differentiation with respect to the new variable x .

To progress we make the assumption

$$g(x) = kh(x) \text{ and } y = \frac{h'}{h}. \quad (14)$$

Then the transformation (14) enables us to write (13) in the form

$$y' + \frac{2a^2bx^{2(b+c)} + acx^{b+c}(2ax^{b+c} - k) + (1-b)c^2k^2 - c^3k^2}{ckx(ck - ax^{b+c})}y + \left(\frac{3ax^{b+c}}{2ck} + \frac{1}{2}\right)y^2 + \frac{a^2b(b+2c)x^{2(b+c-1)}}{2ck(ax^{b+c} - ck)} = 0. \quad (15)$$

Observe that (15) is a Riccati equation in the quantity y . Riccati equations can be transformed to second order linear equations.

We let

$$u(x) = \exp \left[\int^x \left(\frac{3ax^{b+c}}{2ck} + \frac{1}{2} \right) y(x) dx \right]. \quad (16)$$

Note that in equation (16) the term $\left(\frac{3ax^{b+c}}{2ck} + \frac{1}{2} \right)$ is the coefficient of the quantity y^2 in (15). Using (16) equation (15) is transformed to

$$u'' + \gamma(x)u' + \zeta(x)u = 0, \quad (17)$$

where

$$\begin{aligned} \gamma(x) = & \left[6a^3bx^{3(b+c)} + a^2cx^{2(b+c)} \left(6ax^{b+c} + (5b-3)k \right) - c^3k^2 \right. \\ & \left. \left(6ax^{b+c} + (b-1)k \right) + ac^2kx^{b+c} \left(5ax^{b+c} + (2-6b)k \right) \right. \\ & \left. - c^4k^3 \right] \left[ckx \left(-3a^2x^{2(b+c)} + 2ackx^{b+c} + c^2k^2 \right) \right]^{-1}, \quad (18a) \end{aligned}$$

$$\zeta(x) = \frac{a^2b(b+2c)x^{2(b+c-1)}(3ax^{b+c} + ck)}{4c^2k^2(ax^{b+c} - ck)}. \quad (18b)$$

Therefore we have the remarkable feature that the second order nonlinear equation (13) has been transformed to the linear equation (17) via the transformations (14) and (16).

Case $b = -c$

If we set $b = -c$ then equation (17) becomes

$$x^2 u'' + xu' + \frac{a^2(3a + ck)}{4k^2(ck - a)}u = 0, \quad (19)$$

which is a simpler form. It is interesting to note that this case produces the Euler equation (19). We can integrate (19) to give

$$u(x) = \tilde{c}_1 \cosh \left(\frac{a\sqrt{3a + ck}}{2k\sqrt{a - ck}} \log(x) \right) + \tilde{c}_2 \sinh \left(\frac{a\sqrt{3a + ck}}{2k\sqrt{a - ck}} \log(x) \right), \quad (20)$$

where \tilde{c}_1 and \tilde{c}_2 are arbitrary constants of integration. Then from (16) we obtain

$$y(x) = \frac{ac\sqrt{3a + ck}}{\sqrt{a - ck}} \frac{c_1 x^{\frac{a\sqrt{3a + ck}}{k\sqrt{a - ck}}} - c_2}{x(3ax^c + ck) \left(c_1 x^{\frac{a\sqrt{3a + ck}}{k\sqrt{a - ck}}} + c_2 \right)}, \quad (21)$$

where $c_1 = \tilde{c}_1 + \tilde{c}_2$ and $c_2 = \tilde{c}_1 - \tilde{c}_2$.

Hence we have the potentials

$$A = \frac{1}{f(t)} \left(m \left[r^{-1/c} \exp \left(\int^t \frac{dt}{af(t)} \right) \right]^{\frac{a\sqrt{3a+ck}}{2k\sqrt{a-ck}}} + n \left[r^{-1/c} \exp \left(\int^t \frac{dt}{af(t)} \right) \right]^{-\frac{a\sqrt{3a+ck}}{2k\sqrt{a-ck}}} \right)^{\frac{2ck}{3a+ck}}, \quad (22a)$$

$$B = k \frac{f(t)}{r} A, \quad (22b)$$

where $m = c_1 c_3^{\frac{3a+ck}{2ck}}$ and $n = c_2 c_3^{\frac{3a+ck}{2ck}}$ are constants. This is a new solution to the master equation.

Hence we have the potentials

$$A = \frac{1}{f(t)} \left(m \left[r^{-1/c} \exp \left(\int^t \frac{dt}{af(t)} \right) \right]^{\frac{a\sqrt{3a+ck}}{2k\sqrt{a-ck}}} + n \left[r^{-1/c} \exp \left(\int^t \frac{dt}{af(t)} \right) \right]^{-\frac{a\sqrt{3a+ck}}{2k\sqrt{a-ck}}} \right)^{\frac{2ck}{3a+ck}}, \quad (22a)$$

$$B = k \frac{f(t)}{r} A, \quad (22b)$$

where $m = c_1 c_3^{\frac{3a+ck}{2ck}}$ and $n = c_2 c_3^{\frac{3a+ck}{2ck}}$ are constants. This is a new solution to the master equation. The line element for this case is

$$ds^2 = \left[\left(m\psi^{1/2} + n\psi^{-1/2} \right)^{\frac{2ck}{3a+ck}} \right]^2 \left(- \left[\frac{1}{f(t)} \right]^2 dt^2 + \left[\frac{k}{r} \right]^2 [dr^2 + r^2 d\Omega^2] \right), \quad (23)$$

$$\text{where } \psi = \left[r^{-1/c} \exp \left(\int^t \frac{dt}{af(t)} \right) \right]^{\frac{a\sqrt{3a+ck}}{k\sqrt{a-ck}}}.$$

The matter variables become

$$\begin{aligned}
 \mu &= \frac{2(a^2 - ack + c^2 k^2)(ck(m^2 \psi^2 - 4mn\psi + n^2) - 6amn\psi)}{ck^3(a - ck)(3a + ck)(m\psi + n)^2 (\psi^{-1/2}(m\psi + n))^{\frac{4ck}{3a+ck}}}, \\
 p_{\parallel} &= \frac{2a(6amn\psi - ck(m^2 \psi^2 - 4mn\psi + n^2))}{k^2(ck - a)(3a + ck)(m\psi + n)^2 (\psi^{-1/2}(m\psi + n))^{\frac{4ck}{3a+ck}}}, \\
 p_{\perp} &= \frac{(a + ck)(\psi^{-1/2}(m\psi + n))^{-\frac{4ck}{3a+ck}} (12amn\psi + ck(m\psi + n)^2)}{ck^3(3a + ck)(m\psi + n)^2 (\psi^{-1/2}(m\psi + n))^{\frac{4ck}{3a+ck}}}, \\
 q &= \left[\frac{k}{r} \left(m\psi^{1/2} + n\psi^{-1/2} \right)^{\frac{2ck}{3a+ck}} \right]^{-1} p_{\parallel}.
 \end{aligned}$$

From the above we generate the linear barotropic equation of state

$$p_{\parallel} = \lambda \mu, \quad \lambda = \frac{ack}{a^2 - ack + c^2 k^2}, \quad (25)$$

provided that $k \neq -\frac{3a}{c}$.

Case: $k = -3a/c$ and $b = -c$

If we set $k = -3a/c$ and $b = -c$, then equation (13) becomes

$$24x^2hh'' - 24x^2h'^2 + 24xhh' + c^2h^2 = 0, \quad (26)$$

which is greatly simplified. Now (26) can be integrated to give

$$h(x) = \frac{nx^m}{\exp\left[\frac{c^2}{48}\log^2(x)\right]}, \quad (27)$$

where m and n are constants of integration.

Case: $k = -3a/c$ and $b = -c$

If we set $k = -3a/c$ and $b = -c$, then equation (13) becomes

$$24x^2hh'' - 24x^2h'^2 + 24xhh' + c^2h^2 = 0, \quad (26)$$

which is greatly simplified. Now (26) can be integrated to give

$$h(x) = \frac{nx^m}{\exp\left[\frac{c^2}{48}\log^2(x)\right]}, \quad (27)$$

where m and n are constants of integration.

Hence we obtain the metric functions

$$A = \frac{n}{f(t)} \frac{\left[r^{-1/c} \exp\left(\int^t \frac{dt}{af(t)}\right)\right]^m}{\exp\left(\frac{c^2}{48}\log^2\left[r^{-1/c} \exp\left(\int^t \frac{dt}{af(t)}\right)\right]\right)}, \quad (28a)$$

$$B = -\frac{3a}{c} \frac{f(t)}{r} A. \quad (28b)$$

This is also a new solution to the master equation (6).

The line element for this case given by

$$ds^2 = \left[\frac{n\varphi^m}{\exp\left(\frac{c^2}{48} \log^2 \varphi\right)} \right]^2 \left[- \left(\frac{1}{f(t)} \right)^2 dt^2 + 9 \left(\frac{a}{cr} \right)^2 [dr^2 + r^2 d\Omega^2] \right], \quad (29)$$

where $\varphi = r^{-1/c} \exp\left(\int^t \frac{dt}{af(t)}\right)$.

The line element for this case given by

$$ds^2 = \left[\frac{n\varphi^m}{\exp\left(\frac{c^2}{48} \log^2 \varphi\right)} \right]^2 \left[- \left(\frac{1}{f(t)} \right)^2 dt^2 + 9 \left(\frac{a}{cr} \right)^2 [dr^2 + r^2 d\Omega^2] \right], \quad (29)$$

where $\varphi = r^{-1/c} \exp\left(\int^t \frac{dt}{af(t)}\right)$. The matter variables become

$$\begin{aligned} \mu &= \frac{13(c^4 \log^2 \varphi - 48mc^2 \log \varphi + 24(c^2 + 24m^2)) \exp\left(\frac{1}{24}c^2 \log^2 \varphi\right)}{2592a^2 n^2 \varphi^{2m}}, \\ p_{\parallel} &= \frac{(48mc^2 \log \varphi - c^4 \log^2 \varphi - 24(c^2 + 24m^2)) \exp\left(\frac{1}{24}c^2 \log^2 \varphi\right)}{864a^2 n^2 \varphi^{2m}}, \\ p_{\perp} &= \frac{(48mc^2 \log \varphi - c^4 \log^2 \varphi + 48c^2 + 576m^2) \exp\left(\frac{1}{24}c^2 \log^2 \varphi\right)}{648a^2 n^2 \varphi^{2m}}, \\ q &= \left[\frac{3a}{cr} \frac{n\varphi^m}{\exp\left(\frac{c^2}{48} \log^2 \varphi\right)} \right]^{-1} p_{\parallel}. \end{aligned}$$

We also have $p_{\parallel} = \lambda\mu$, $\lambda = -\frac{3}{13}$, which is linear and barotropic.

Case: $b = \left(\pm \frac{\sqrt{3}}{3} - 1 \right) c$

In this case equation (13) becomes

$$6gg'h' - 3(2gg'' + g'^2)h \pm 2\sqrt{3}g'h^2 = 0. \quad (31)$$

The advantage of (31) is that it is a Bernoulli equation in h . Here g is unspecified. We integrate (31) to obtain

$$h(t) = \pm \frac{\sqrt{3}}{2} \frac{g' \sqrt{g}}{\sqrt{g} + d}, \quad (32)$$

where d is a constant of integration.

Case: $b = \left(\pm \frac{\sqrt{3}}{3} - 1 \right) c$

In this case equation (13) becomes

$$6gg'h' - 3(2gg'' + g'^2)h \pm 2\sqrt{3}g'h^2 = 0. \quad (31)$$

The advantage of (31) is that it is a Bernoulli equation in h . Here g is unspecified. We integrate (31) to obtain

$$h(t) = \pm \frac{\sqrt{3}}{2} \frac{g' \sqrt{g}}{\sqrt{g} + d}, \quad (32)$$

where d is a constant of integration. Hence the potentials functions become

$$A = \pm \frac{\sqrt{3}}{2} \frac{g' \sqrt{g}}{\sqrt{g} + d} r^{\pm \frac{\sqrt{3}}{3}}, \quad (33a)$$

$$B = gr^{\pm \frac{\sqrt{3}}{3} - 1}, \quad (33b)$$

which is a new exact solution for the shear-free model.

The line element for this model becomes

$$ds^2 = -\frac{3}{4}g'^2 r^{\pm 2\frac{\sqrt{3}}{3}} dt^2 + g^2 r^{\pm \frac{2\sqrt{3}}{3}-2} [dr^2 + r^2 d\Omega^2] . \quad (34)$$

from (33). In the above we have set the arbitrary constant $d = 0$ without any loss of any generality.

The line element for this model becomes

$$ds^2 = -\frac{3}{4}g'^2 r^{\pm 2\frac{\sqrt{3}}{3}} dt^2 + g^2 r^{\pm \frac{2\sqrt{3}}{3}-2} [dr^2 + r^2 d\Omega^2]. \quad (34)$$

from (33). In the above we have set the arbitrary constant $d = 0$ without any loss of any generality. The matter variables become

$$\begin{aligned} \mu &= \frac{14r^{\mp \frac{2}{\sqrt{3}}}}{3g^2}, \\ p_{\parallel} &= -\frac{4r^{\mp \frac{2}{\sqrt{3}}}}{3g^2}, \\ p_{\perp} &= -\frac{r^{\mp \frac{2}{\sqrt{3}}}}{g^2}, \\ q &= \left[gr^{\pm \frac{\sqrt{3}}{3}-1}\right]^{-1} p_{\parallel}. \end{aligned}$$

This solution also satisfies the barotropic equation of state.

$$p_{\parallel} = \lambda \mu, \quad \lambda = -\frac{2}{7}, \quad (36)$$

which is linear.

Summary

- We developed the junction condition equation that relates the radial pressure to the heat flux which is a highly nonlinear partial differential equation in the metric functions.
- We demonstrated that this equation admits three Lie point symmetries.
- Using the general linear combination of these symmetries we reduced the governing highly nonlinear partial differential equation to ordinary differential equations.
- By solving the reduced ordinary differential equations and transforming to the original variables we obtained exact solutions for the master equation.
- We present the line element explicitly in each case and show that our solutions obey the linear barotropic equation of state.

References



Abebe G Z, Govinder K S and Maharaj S D,
Lie symmetries for a conformally flat radiating star,
Int. J. Theor. Phys. **52**, 3244 (2013).



Abebe G Z, Maharaj S D and Govinder K S,
Geodesic models generated by Lie symmetries,
Gen. Relativ. Gravit. **46**, 1650 (2014a).



Abebe G Z, Maharaj S D and Govinder K S,
Generalized Euclidean stars with equation of state,
Gen. Relativ. Gravit. **46**, 1733 (2014b).



Abebe G Z, Maharaj S D and Govinder K S,
A group theoretic approach to shear-free radiating stars,
Eur. Phys. J. C, **75**, 496 (2015).



Govinder K S and Govender M,
A general class of Euclidean stars,
Gen. Relativ. Gravit. **44**, 147 (2012).

Thank You!